

# Keep Your Cells Happy

Eppendorf Bioreactor Systems meet the high demands of stem cells perfectly. Parallel optimization testing, precise control, around the clock monitoring and comprehensive options allow for the most effective process. Eppendorf Bioreactor Systems are proven for various ES, MSC and iPS cell lines. Seamlessly transition your stem cell experiments to the next level with the DASBox® Mini.

View the DASBox® Mini at  
[www.eppendorf.com/DASBox](http://www.eppendorf.com/DASBox)

# Wild-type MeCp2 expression coincides with age-dependent sensory phenotypes in a female mouse model for Rett syndrome

Michael Mykins  | Dana Layo-Carris  | Logan Reid Dunn  | David Wilson Skinner |  
Alexandra Hart McBryar | Sarah Perez | Trinity Rose Shultz | Andrew Willems  |  
Billy You Bun Lau  | Tian Hong  | Keerthi Krishnan 

Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, Tennessee, USA

## Correspondence

Keerthi Krishnan, Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, TN, USA.  
Email: [krishnan@utk.edu](mailto:krishnan@utk.edu)

## Funding information

International Rett Syndrome Foundation; National Institute of Mental Health, Grant/Award Number: R15MH124042; University of Tennessee, Knoxville; National Science Foundation Graduate Research Fellowship; National Institute of General Medical Sciences of the National Institutes of Health, Grant/Award Number: R01GM140462

## Abstract

Rett syndrome is characterized by an early period of typical development and then, regression of learned motor and speech skills in girls. Loss of MECP2 protein is thought to cause Rett syndrome phenotypes. The specific underlying mechanisms from typical developmental trajectory to regression features throughout life are unclear. Lack of established timelines to study the molecular, cellular, and behavioral features of regression in female mouse models is a major contributing factor. Due to random X-chromosome inactivation, female patients with Rett syndrome and female mouse models for Rett syndrome (*Mecp2*<sup>Heterozygous</sup>, Het) express a functional copy of wild-type MeCp-2 protein in approximately half of all cells. As MeCp2 expression is regulated during early postnatal development and experience, we characterized the expression of wild-type MeCp2 in the primary somatosensory cortex of female Het mice. Here, we report increased MeCp2 levels in non-parvalbumin-positive neurons of 6-week-old adolescent Het relative to age-matched wild-type controls, while also displaying typical levels of perineuronal net expression in the barrel field subregion of the primary somatosensory cortex, mild tactile sensory perception deficits, and efficient pup retrieval behavior. In contrast, 12-week-old adult Het express MeCp2 at levels similar to age-matched wild-type mice, show increased perineuronal net expression in the cortex, and display significant tactile sensory perception deficits. Thus, we have identified a set of behavioral metrics and the cellular substrates to study regression during a specific time in the female Het mouse model, which coincides with changes in wild-type MeCp2 expression. We speculate that the precocious increase in MeCp2 expression within specific cell types of adolescent Het may provide compensatory benefits at the behavioral level, while the inability to further increase MeCp2 levels leads to regressive behavioral phenotypes over time.

Michael Mykins, Dana Layo-Carris, and Logan Reid Dunn are co-first authors.

Edited by Junie Paula Warrington and Samantha McLean. Reviewed by Zhaolan Zhou.

## KEYWORDS

adolescence and adulthood, barrel cortex, female mice, tactile sensation, X-linked disorders, maternal behavior, MeCp2, mosaicism, perineuronal nets, Rett syndrome

## 1 | INTRODUCTION

Rett syndrome (RTT) is a neuropsychiatric disorder, predominantly caused by mutations in the X-linked gene Methyl CpG-binding protein 2 (*MECP2*) (Amir et al., 1999; Hagberg et al., 1983; Rett, 1966). Most surviving patients with RTT are females who are heterozygous for *MECP2* mutations. Diagnostic criteria for RTT include partial or complete loss of acquired purposeful motor skills or spoken language during a regression period in the first few years of life. Additionally, primary sensory processing is disrupted in RTT patients (Buchanan et al., 2019; Djukic et al., 2012; Djukic & McDermott, 2012; Key et al., 2019; LeBlanc et al., 2015; Merbler et al., 2020; Nomura, 2005; Peters et al., 2015; Stallworth et al., 2019; Symons et al., 2019). Such disruptions during development and adulthood contribute to motor dysfunction and social communication deficits in girls and women with RTT. Patients typically survive into middle age with periods of plateau and regression throughout life (Bebbington et al., 2008; Buchanan et al., 2019; Charman et al., 2002; Djukic et al., 2012; Djukic & McDermott, 2012; Einspieler & Marschik, 2019; Kerr, 1995; Kirby et al., 2010; Nomura, 2005). Rodent models recapitulate many of the features of RTT, including sensory processing, social communication deficits, and motor difficulties (Durand et al., 2012; Goffin et al., 2012; Ito-Ishida et al., 2015; Krishnan et al., 2015, 2017; Lau, Krishnan, et al., 2020; Lee et al., 2017; Lo et al., 2016; Orefice et al., 2016; Su et al., 2015). Although much is known about the characteristics of RTT phenotypes, the underlying cause for regression during various stages of life and the changes in syndromic phenotypes over time are unclear. MeCp2 is a master gene regulator that “reads” DNA methylation signals, acts as a scaffolding protein for activator or repressor protein complexes, and/or regulates chromatin conformation and 3D genome organization (Boxer et al., 2020; Chahrour et al., 2008; Ito-Ishida et al., 2020; Kruusvee et al., 2017; Lewis et al., 1992; Linhoff et al., 2015; Lyst et al., 2013; Meehan et al., 1992). In neurons, MeCp2 is expressed at a level equivalent to that of core histones (Skene et al., 2010). The prevailing idea is that loss of MeCp2 expression and function is the major driver for RTT phenotypes; however, the role of the wild-type MeCp-2 protein in half of the cells in RTT is unknown.

In RTT, random X-chromosome inactivation leads to mosaic wild-type MeCp2 expression in approximately 50% of most major cell types in the body, including neurons (Samaco et al., 2013; Young & Zoghbi, 2004). Due to this inherent mosaicism, individual girls and women with RTT, as well as individual female heterozygous mice (*Mecp2*<sup>heterozygous</sup>, Het), are unique. However, the impact of mosaic MeCp2 expression on cellular function in Het is not fully understood. We hypothesize that the mosaicism in

## Significance

Although in its early stages, we believe these findings provide a novel path to tackle neurobiological underpinnings of X-linked neurodevelopmental disorders. The path focuses on the regulation of wild-type protein expression in approximately half of the cells and homeostatic plasticity in the neural networks with non-expressing cells in females over time and age. Ultimately, this observation opens up new avenues to pursue therapeutic strategies to treat Rett syndrome. The impact of these observations will be far-reaching as this well-reported neuropsychiatric disorder typically serves as a model for monogenic, yet complex social and sensory disorders.

Het contributes to the observed syndromic phenotypes, due to likely compensation and adaptations in individual cells throughout life. In rodents, MeCp2 expression increases postnatally, concomitant with the emergence of appropriate critical periods, and is generally thought to be stable in adolescence and adulthood (Braunschweig et al., 2004; Krishnan et al., 2015; Picard & Fagiolini, 2019; Skene et al., 2010; Yagasaki et al., 2018). MeCp2 is variably expressed across cell types, with higher expression in GABAergic interneurons in comparison to pyramidal neurons within the same region (Braunschweig et al., 2004; Ito-Ishida et al., 2015; Skene et al., 2010; Takei et al., 2021; Yagasaki et al., 2018). Intriguingly, the status of wild-type MeCp2 expression over time, in Het, is unknown. Due to its significant roles in gene regulation (Chahrour et al., 2008; Chen et al., 2015; Fuks et al., 2003; Gabel et al., 2015; Johnson et al., 2017; Jones et al., 1998; Klose et al., 2005; Lyst & Bird, 2015; Tao et al., 2009; Zhou et al., 2006) and the timing of experience-dependent plasticity (Durand et al., 2012; Krishnan et al., 2015, 2017; Lau, Krishnan, et al., 2020; Morello et al., 2018; Noutel et al., 2011; Patrizi et al., 2020; Picard & Fagiolini, 2019), investigation of wild-type MeCp2 expression in Het across ages is essential to unravel the mechanisms connecting genotype to phenotype throughout the lifetime of the individual.

Initial characterization of *Mecp2*-heterozygous females using standard behavioral assays classified the 6-week-old adolescent Het as pre-symptomatic, whereas 12-week-old adult females would be considered at the early symptomatic stage (Samaco et al., 2013). Previously, we established the pup retrieval task during ethologically relevant alloparental behavior, as a model to specifically address the cellular and neural circuitry basis for social communication and complex sensorimotor integration

(Krishnan et al., 2017; Lau, Krishnan, et al., 2020; Stevenson et al., 2021). This innate task in mice involves females retrieving scattered pups back to the nest (Alsina-Llanes et al., 2015; Beach & Jaynes, 1956; Champagne et al., 2007; Champagne & Curley, 2016; Hemel, 1973; Koch & Ehret, 1989; Lonstein et al., 2015; Stern, 1996; Stern & Mackinnon, 1978). Adult Het are inefficient at the pup retrieval compared to age-matched wild-type (WT) females (Krishnan et al., 2017; Stevenson et al., 2021). Furthermore, adult Het exhibit atypical tactile interactions with pups involving the whiskers and face, with decreased grooming behavior and increased proximal pup approaches (Stevenson et al., 2021). Together, these results suggest abnormal tactile processing in Het, which could contribute to their inefficient pup retrieval. Tactile processing is particularly relevant to alloparental care, as sensory cues from offspring are singularly essential for the early retention and maintenance of maternal behavior (Hinde & Davies, 1972; Lonstein & Stern, 1999; Maier, 1963; Morgan et al., 1992; Rosenblatt, 1967; Stern, 1996; Stern & Kolonie, 1989; Stern & Mackinnon, 1978). Thus, we focused on the tactile sensory perception modality in adolescent and adult Het using texture discrimination and object recognition tasks (Antunes & Biala, 2012; Diamond et al., 2008; Orefice et al., 2016, 2019; Wu et al., 2013).

Here, we report the first evidence of a precocious increase of wild-type MeCp2 expression within MeCp2-expressing cells of 6-week-old adolescent Het relative to age-matched WT; these MeCp2 expression changes are observed in non-parvalbumin positive (PV+) GABAergic neurons, which exhibit lower levels of MeCp2, but not in PV+ GABAergic neurons exhibiting higher MeCp2 levels. This cell-type specific expression difference is not maintained in 12-week-old adult Het, suggesting that cells expressing wild-type MeCp-2 protein in Het are regulated by age-specific mechanisms. This MeCp2 expression phenotype in adolescent Het is concomitant with typical structural plasticity, as measured by the expression of specialized extracellular matrix structures ensheathing PV neurons called perineuronal nets (PNNs). Furthermore, systematic, frame-by-frame analyses of texture and object investigation tasks in conjunction with pup retrieval behavior reveal that adolescent Het behave similar to adolescent WT in both simple tactile and complex social behaviors. Adult Het mice, however, are known to be inefficient at pup retrieval (Krishnan et al., 2017; Stevenson et al., 2021) and display overt tactile sensory perception issues, suggesting that observed tactile hyposensitivity in Het is age-related. Based on these findings, we hypothesize that precocious increase of wild-type MeCp2 expression in specific cell types of adolescent Het provides compensatory benefits at the behavioral level, while the inability to further increase MeCp2 levels with age leads to observed RTT stereotypies over time. Together, these novel results open new directions of exploration on the long-term, cell-type-specific roles of MeCp2 expression, regulation, and function in Het while emphasizing the need to analyze dynamics of cellular and behavioral phenotypes.

## 2 | MATERIALS AND METHODS

### 2.1 | Animals

We used the following mouse strains: CBA/CaJ (RRID:IMSR\_JAX:000654) and *Mecp2*<sup>heterozygous</sup> (Het) (C57BL/6J, B6.129P2(C)-*Mecp2*<sup>tm1.1Bird</sup>/J, RRID:IMSR\_JAX:003890) (Guy et al., 2001). All animals were group-housed by sex after weaning, raised on a 12/12-h light/dark cycle (lights on at 7 am), and received food and water ad libitum. Behavioral experiments were performed using Het and female *Mecp2*<sup>WT</sup> siblings (WT) at either 6 (adolescent) or 10–12 (adult) weeks of age between the hours of 9 am and 6 pm (during the light cycle). All procedures were conducted in accordance with the National Institutes of Health's Guide for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee at the University of Tennessee-Knoxville.

### 2.2 | Immunohistochemistry

All animals were anesthetized with isoflurane and perfused with 4% paraformaldehyde (PFA) in 1× Phosphate Buffer Saline (PBS) immediately after behavior. Extracted brains were stored in 4% PFA overnight at 4°C before storing long term in PBS. Tissues were processed in cohorts (one WT and one Het) as previously described (Lau, Layo, et al., 2020). Briefly, coronal brain sections (100µm thickness) encompassing the entire primary somatosensory cortex (SS1) were collected using a freezing microtome. We denoted the left hemisphere by poking a small hole in the ventral part of the brain. Free-floating sections were blocked with a 10% normal goat serum and .5% Triton X-100 in 1× PBS. Blocking solution at half the original concentration was used for antibody incubations. Tissues were incubated with biotin-conjugated WFA lectin (1:500; Sigma-Aldrich Cat# L-1516, RRID:AB\_2620171), MeCp2 primary antibody (1:1000; Cell Signaling Technology Cat# 3456, RRID:AB\_2143849), and PV primary antibody (1:1000; Sigma-Aldrich Cat# P3088, RRID:AB\_477329) followed by secondary antibodies conjugated with AlexaFluor-488 (Molecular Probes Cat# S32354, RRID:AB\_2315383), Texas-Red (Molecular Probes Cat# T-6391, RRID:AB\_2556779), and AlexaFluor-405 (Molecular Probes Cat# A48255, RRID:AB\_2890536) (all at 1:1000) to label PNNs, MeCp2, and PV (if applicable), respectively. Tissues were counterstained with DAPI before mounting section onto slides.

### 2.3 | Microscopy and image analysis

#### 2.3.1 | Image acquisition

##### 2.3.1.1 | *MeCp2*

Image acquisition of both MeCp2 and DAPI signals was performed using a VT-Hawk 2D-array scanning confocal microscope (Visitech Intl.) mounted to an IX-83 inverted optical microscope (Olympus; RRID:SCR\_020344). Z-series images with a step interval of .5 µm

were collected using a 60× objective lens (Nikon) in MetaMorph (Molecular Devices; [RRID:SCR\\_002368](#)). Settings for each laser channel were determined by using the histogram function in MetaMorph. Acquisition settings were determined using the top 25.0 μm of WT tissues, and these settings were used to image all conditions in the 16-bit range. PV and PNN images ([Figure 1a](#)) were acquired in the same manner using a 40× objective lens (Nikon). Images were collected from the primary somatosensory cortical layer IV of barrel field subregion (S1BF) in both hemispheres of two sections per brain.

#### 2.3.1.2 | PNN

Single-plane images of coronal tissue sections were collected using a Keyence epifluorescence microscope (Keyence BZ-X710; [RRID:SCR\\_017202](#)) with a 10× objective and stitched using the BZ-X Analyzer software. We determined exposure settings for each cohort based on Het PNN intensity, as we previously showed atypical increased expression of PNNs in adult Het (Lau, Layo, et al., 2020). To reduce overexposure, we set the exposure time to the point when the first saturated pixel appeared in the sample, then reduced the exposure time by one unit. This procedure was performed in both hemispheres for all Het sections, and the lowest exposure time occurring most frequently was used to image all sections within that cohort.

### 2.3.2 | Image processing & analysis

#### 2.3.2.1 | MeCp2

Nuclear expression of MeCp2 was quantified using ImageJ ([RRID:SCR\\_003070](#)) (Schneider et al., 2012). For initial background subtraction, a duplicate of the original image was created and processed by applying a median filter with a 32-pixel radius. The filtered image was then subtracted from the original image using the Image Calculator tool to produce 16-bit images for quantification. Mean MeCp2 intensity was quantified within individual nuclei by creating 3.0–5.0 μm maximum intensity projections (16-bit grey-scale) within the top 25.0 μm of each tissue. Projections were generated using the same image plane for both the DAPI and MeCp2 channels and utilized for region-of-interest (ROI) generation via the Polygon drawing tool and quantification, respectively. DAPI signal, PNN presence or absence, and MeCp2 expression status were identified for individual cells using merged tricolor images. Statistical analyses and figures were generated using GraphPad Prism ([RRID:SCR\\_002798](#)). The percentage of nuclei expressing MeCp2 was calculated using the EzColocalization ImageJ plugin (Stauffer et al., 2018). Briefly, the DAPI and MeCp2 z-series image stacks were used for reporter channels 1 and 2, respectively, with default threshold algorithms applied. The mean Mander's colocalization coefficient (MCC) for MeCp2 colocalization with DAPI signal was collected across all imaging planes. This process was repeated for the four images collected for each animal (2 per hemisphere) and averaged to yield an animal-specific mean MCC value. To

compare MCC values in a standardized manner, the average MCC value for all WT images across both ages was calculated. Then, the mean MCC value for each WT and Het replicate was divided by the WT average MCC and multiplied by 100 to approximate the percentage of nuclei expressing MeCp2.

#### 2.3.2.2 | PNN

Images were mapped and ROIs for S1BF were outlined by overlaying the images with coronal maps from Paxinos and Franklin's mouse atlas (4th edition, 2013). The density of high-intensity PNNs was quantified as previously described (Lau, Layo, et al., 2020) using ImageJ. Briefly, high-intensity PNNs were quantified by maximizing the contrast of each image and manually marking the signals that were 80% or more of the original PNN. Density was calculated as total number of high-intensity PNNs divided by ROI area. We also analyzed the full spectrum of PNN expression in the mapped images. Eight-bit histograms specific to the green channel (PNNs) were generated using the Histogram tool in ImageJ. We extracted the distributions of strong PNN expression from all images using two-component Gaussian mixture models (GMM; see [Figure 2](#)) (Bibbings et al., 2019). The means of these distributions (red traces in [Figure 2d](#)) were used as representative summary statistics in subsequent analyses. We calculated intraclass correlation coefficients (ICCs) to show that the ratio of inter-individual variance to the total variance was moderate (.218), such that comparisons on genotypes can be made. We built a linear mixed-effects model to confirm that the inter-individual variability did alter the statistical significance (Yu et al., 2022). The increased variability in Het compared to WT is likely due to natural variation due to random X-chromosome inactivation patterns in Het.

### 2.4 | Pup retrieval task

#### 2.4.1 | Behavior paradigm

The pup retrieval task was performed as previously described (Stevenson et al., 2021). Two 4.5-week-old naïve females (one WT and one Het) with no prior experience of pup retrieval were co-housed with a pregnant CBA/CaJ female (10–12 weeks of age) 3–5 days before birth; naïve females were termed “surrogates” upon cohousing. Once the pups were born (postnatal day 0; P0), we performed the pup retrieval task with the surrogates for six consecutive days. The behavior was performed in the home cage placed inside of a sound- and light-proof box as follows: One surrogate was placed in the home cage with three to five pups (*habituation*; 5 min), the pups were removed from the nest (*isolation*; 2 min), and the pups were scattered throughout the home cage by placing them in the corners and center, allowing the surrogate to retrieve pups back to the nest (*retrieval*; 5 min). The nest was left empty if there were fewer than five pups. The assay was performed in the dark and recorded using infrared cameras during all phases of each trial. If all pups were

not retrieved to the nest by the 5-min mark of the retrieval phase, we placed pups back in the nest in preparation for the next surrogate. After the behavior was completed, all mice were returned to the home cage.

## 2.4.2 | Behavior analysis

All videos were coded, so the analyzers were blind to the identity of the mice. Each video was manually scored using a latency index (the amount of time to retrieve all pups back to the nest out of 5 min) and the number of errors (surrogate-pup interactions that did not result in successful retrieval). Latency index was calculated as:

$$\text{Latency index} = [(t_1 - t_0) + (t_2 - t_0) + \dots + (t_n - t_0)] / (n \times L)$$

where  $n$  = number of pups outside the nest,  $t_0$  = start time of trial (s),  $t_n$  = time of the  $n$ th pup's successful retrieval to the nest (s),  $L$  = total trial length (300s). Statistical analyses and figures were generated using GraphPad Prism.

## 2.5 | Tactile perception tasks

WT and Het underwent a 5-day battery of behavioral tests to assay tactile processing through the whiskers (Figure 4). The behavior chamber used for tactile assays was a 40×40×40cm plexiglass box with two sets of holes in the center of the plexiglass floor for the texture/object inserts. Holes were drilled to be equidistant from the center, with a 4.0-cm spacing between hole sets. Texture discrimination plexiglass inserts were .5cm (thickness)×4.0cm (width)×15.0cm (height). Rodents use their whiskers to both detect and discriminate between textures and objects (Rodgers et al., 2021). We chose to test the ability of experimental mice to discriminate between textures using both large and small textural differences. Sandpaper textures were designed to slide on and off the inserts to be cleaned between animals. The textures were made of 3M sandpaper of varying grit: a course 60-grit, a medium 120-grit, and a fine 220-grit. Inserts used for novel object recognition were plexiglass spheres (4.0 cm diameter) and plexiglass cubes (64cm<sup>3</sup>) (Orefice et al., 2016; Wu et al., 2013). The chamber and inserts were cleaned with MB-10 chlorine dioxide disinfectant and allowed to fully dry before the start of each behavior, and an additional 5-min period was allotted between tests to eliminate scent biases. All behaviors were performed in the dark and recorded with an infrared camera.

### 2.5.1 | Behavior paradigm

#### 2.5.1.1 | Habituation

On days 1 and 2, mice were placed individually into the behavior chamber for 10 min to habituate themselves with the new environment.

#### 2.5.1.2 | Texture discrimination

On days 3 and 4, texture discrimination assays began with a familiarization trial. Experimental mice were individually placed into the chamber with two identical texture inserts and allowed to investigate for 10min. Afterward, the mouse was returned to the home cage for 5min. During this time, textures and inserts for familiarization were replaced with new sets of inserts and textures for use in the testing trial. After 5min, the mouse was returned to the behavior chamber for a second 10-min trial (testing phase). Familiarization phases of each behavior were performed using two course texture inserts. During testing phase on days 3 and 4, we replaced one insert with a medium and fine texture insert, respectively.

#### 2.5.1.3 | Novel object recognition

On day 5, a novel object recognition assay was performed similar to the texture discrimination assay with minor changes. The familiarization trial was performed using two plexiglass cubes. For the testing trial, one plexiglass cube was replaced by a plexiglass sphere.


### 2.5.2 | Behavior analysis

Behaviors for both texture and object investigation were manually scored using Datavyu software (RRID:SCR\_003587; Datavyu Team, version 1.3.7, 2014). All analyzers were blind to the age and genotype of the mice. The videos were scored for the time intervals in which the mouse interacted with the novel and familiar textures/objects throughout the 10-min testing trial. We classified interactions into three categories, as shown in Table 1, based on pilot observations and previous studies (Orefice et al., 2016; Pacchiarini et al., 2017). We excluded mice from analysis if the investigation time of textures/objects during a given trial did not exceed 2s (Orefice et al., 2016; Wu & Dyck, 2018) (Table 2). After manual scoring, interaction sequences for each animal were extracted from Datavyu for quantitative analyses. We classified number of contacts, total investigation time, and duration per interactions with textures/objects into whisker, forepaw, and whole-body climbing. Our results are reported as raw number of contacts, raw total investigation times, and raw duration per interaction split by interaction type, age, and genotype. We conducted unpaired Mann-Whitney  $U$  tests to determine statistical significance between genotypes and across ages within the same genotype. Statistical analysis and data visualization were performed in R studio (RRID:SCR\_000432).

## 3 | RESULTS

### 3.1 | Wild-type MeCp2 expression is dynamically regulated in Het over age and cell types

MeCp-2 protein expression varies in different cortical cell types (Ito-Ishida et al., 2015; Takei et al., 2021). However, it is currently unknown if cell-type-specific MeCp2 expression is stable over age



**FIGURE 1** MeCp2 is expressed differentially between cell types in S1BF of adolescent and adult WT and Het. (a) Representative 40x confocal images from Layer IV S1BF showing PNN colocalization with PV in MeCp2+ nuclei. Arrows (non-PV) and arrowheads (PV) point to representative nuclei for quantification. Scale bar = 10 μm. (b) Mean intensity of MeCp-2 protein from PV and non-PV nuclei is represented as a fold difference (PV: non-PV ratio; number of replicates same as reported for figure f, g). (c, d) Top row: Representative confocal images of nuclei stained with DAPI and MeCp2 from non-PV (PNN-negative cells) (c) and PNN-positive parvalbumin (PV) interneurons (d) in adolescent and adult naïve WT and Het. Bottom row: Respective maximum intensity projection images of top row, to highlight the differential MeCp2 expression across ages and genotypes. MeCp2 intensity is higher within PV nuclei across ages, and non-PV nuclei of adolescent (6 wk) Het appear more intense compared to age-matched WT or adult (12 wk) Het. Scale bars = 10 μm. (e) Het express MeCp2 in approximately 50% of nuclei, compared to WT across ages, as measured by percentage of MeCp2-positive nuclei in adolescent and adult WT and Het ( $n=2$  images per hemisphere per brain). (f, g) Violin plots of the mean intensity of nuclear MeCp2 within non-PV (f) and PV (g) nuclei. MeCp2 expression increased significantly with age in non-PV nuclei of WT, while MeCp2 expression in non-PV cells of Het remained consistent across ages. Compared to age-matched WT control, 6-week-old Het showed a significant increase in MeCp2 expression in non-PV cells, while 12-week-old Het did not. In PV nuclei, MeCp2 expression increased significantly with age within both genotypes; there were no significant differences between WT and Het MeCp2 expression at either age. Respective ages are indicated on the x-axes, and the number of nuclei from which data were collected is shown in parentheses for each condition ( $n=5$  animals per genotype at adolescent age;  $n=4$  animals per genotype at adulthood). Kruskal–Wallis followed by uncorrected Dunn's test: ns, not significant; \*\* $p < .01$ , \*\*\* $p < .001$ , \*\*\*\* $p < .0001$ . (b, e) Mean  $\pm$  SEM are shown.

or if wild-type MeCp2 expression in Het is similar to age-matched WT controls. Het expression of wild-type MeCp2 was reported to be decreased relative to age-matched WT females; however, the brain region(s) and the specific ages of mice used in this investigation were not reported (Braunschweig et al., 2004). Thus, we performed

immunofluorescent staining and confocal microscopy to first characterize MeCp2 expression within individual nuclei of naïve WT and Het female mice during adolescence and adulthood. We chose to investigate expression within Layer IV of the barrel field subregion of the primary somatosensory cortex (S1BF), the cortical input

layer from the whiskers, due to its importance in tactile sensory perception.


Previously, we reported that adult Het exhibit dysregulation of PV+ GABAergic neurons in the auditory and primary somatosensory cortices (Krishnan et al., 2017; Lau, Krishnan, et al., 2020; Lau, Layo, et al., 2020). Particularly, adult Het mice exhibit increased and atypical expression of perineuronal nets (PNNs) in the S1BF (Lau, Layo, et al., 2020), extracellular matrix structures mainly surrounding cortical PV+ GABAergic interneurons to structurally restrict plasticity (de Vivo et al., 2013; Hartig et al., 1992; Krishnan et al., 2017; Pizzorusso et al., 2002; Ueno et al., 2018). Thus, we focused MeCp2 expression analysis in PNN+ PV nuclei and neighboring PNN-negative nuclei (termed non-PV) within S1BF (Figure 1a).

PNN+ PV nuclei of WT exhibited a roughly threefold higher level of MeCp2 than in non-PV nuclei in Layer IV S1BF across adolescence and adulthood (Figure 1a,b). This fold increase is higher than the 1.25-fold increase reported in Layers II/III (Ito-Ishida et al., 2015). We observed significant increases in MeCp2 expression between adolescence and adulthood in WT females across cell types (Figure 1c,d,f,g). In Het, approximately 50% of the nuclei

(identified by DAPI staining) expressed wild-type MeCp-2 protein at both ages (Figure 1e), in agreement with previous reports (Samaco et al., 2013; Young & Zoghbi, 2004). Strikingly, adolescent Het non-PV nuclei expressed significantly higher levels of MeCp2 than age-matched WT, while adult Het non-PV nuclei did not exhibit such a difference (Figure 1b,c,f). In PV nuclei, MeCp2 expression was significantly increased between adolescence and adulthood in both WT and Het (Figure 1d,g), without significant differences between genotypes within ages. Together, these results show that Het displays a precocious increase of wild-type MeCp2 levels, as well as an age-dependent failure to further increase wild-type MeCp2 expression levels on par with WT females, within specific cell types.

### 3.2 | No changes in perineuronal net expression in S1BF of adolescent Het

MeCp2 regulates the expression of perineuronal nets (PNNs), specialized extracellular matrix structures that structurally restrict changes in synaptic plasticity (Carstens et al., 2016, 2021;



**FIGURE 2** Adolescent Het show similar PNN density as WT within S1BF. (a) Representative epifluorescent 10 $\times$  micrographs of PNNs in S1BF of adolescent WT and Het. Arrows indicate high-intensity PNNs quantified for analysis. (b) Schema of coronal mouse brain sections, depicting rostral (Bregma: .37 mm) to caudal (Bregma: -1.91 mm) regions of S1BF (boxes) analyzed for this study. Coordinate maps are based on Paxinos and Franklin's mouse brain atlas, 4th edition. (c) Statistical analysis of manually counted high-intensity PNNs revealed that the density is similar between adolescent WT and Het. Each circle represents the average high-intensity PNN density of each brain, superimposed on violin plots showing median with distribution of all sections. Individual colors represent cohorts processed and analyzed together. WT:  $n=307$  images, 6 animals. Het:  $n=313$  images, 6 animals. Mann-Whitney test: ns, not significant. (d) All intensity analysis of PNN expression. The top two panels show two distributions of PNN intensity obtained with a two-component Gaussian mixture model for each image (see Materials and Methods). The distributions with higher means (red traces) were used for subsequent analyses, while background signal (blue traces) was excluded. Each trace represents PNN intensity distribution of an image. Insets show histograms of raw intensities for pixels of all images (WT: 135 images and Het: 140 images). Bottom panel: Statistical analysis revealed no significant difference in all PNN intensity between adolescent WT and Het. The graph shows the distribution of the means (small red dots and violin plots) of the red traces in top panels. Large dots show aggregated mean of the red distributions for each color-coded cohort. Box plots within violin plots show 25th, 50th, and 75th quartiles. Linear mixed effect model and  $t$ -test:  $p > .05$ ; ns, not significant.

Donato et al., 2013; Durand et al., 2012; Gogolla et al., 2009; Hou et al., 2017; Kosaka & Heizmann, 1989; Krishnan et al., 2015, 2017; Morello et al., 2018; Nakagawa et al., 1986; Patrizi et al., 2020; Pizzorusso et al., 2002; Sigal et al., 2019; Thompson et al., 2018; Ueno et al., 2017). Adult Het show atypical increases in the expression of mature, high-intensity PNNs in S1BF and the auditory cortex (Krishnan et al., 2017; Lau, Layo, et al., 2020). PNNs are formed on PV neurons from an array of secreted proteins sourced from multiple cell types (Bignami et al., 1992; Carulli et al., 2006, 2007; Kwok et al., 2010; Miyata & Kitagawa, 2017). Thus, we hypothesized that the increased MeCp2 levels within non-PV cells would regulate adolescent Het PNN expression to typical WT levels in a non-cell-autonomous manner.

As we found nuanced changes in lateromedial axis and a hemisphere-specific difference in PNN expression in adult females (Lau, Layo, et al., 2020), we used serial sectioning, immunofluorescent staining, and epifluorescence microscopy to manually quantify high-intensity PNN expression across the entire S1BF (Figure 2a,b). We found no significant difference in high-intensity PNN density between adolescent WT and Het across the entirety of the S1BF (Figure 2c). Additionally, we did not find differences in PNN densities between hemispheres of either adolescent genotype. (WT-L =  $12.845 \pm 1.607$  # PNNs/mm<sup>2</sup>, WT-R =  $18.315 \pm 1.016$ ,  $n = 307$  images, 6 animals; Het-L =  $13.241 \pm .380$ , Het-R =  $14.932 \pm .347$ ,  $n = 313$  images, 6 animals; unpaired Mann-Whitney test).

Differing intensities of PNNs are thought to represent levels of maturity of that PV network in early developing postnatal cortex (Krishnan et al., 2015; Orlando et al., 2012; Pizzorusso et al., 2002; Ye & Miao, 2013). However, it is unclear if the same standards hold true for adolescent cortices. Thus, we performed additional analyses using an automated, unbiased approach to select all pixels that represent PNN expression (Figure 2d, red distributions in top two panels; see Materials and Methods) over background (Figure 2d, blue distributions in top two panels). We observed that PNN expression

in all tissues had multimodal pixel intensity distributions, which consistently included a bell-shaped distribution at higher intensities (Figure 2d insets). This distribution represents specific cortical layers with strong PNN expression (Layers II/III, IV, and V) of interest for quantification. Models consistently captured the distributions of PNN expression in the expected intensity region (Figure 2d, red). By analyzing the mean distribution of these red traces, we observed no significant differences in PNN signal between adolescent WT and Het using a linear mixed-effects model with cohort as a variable (Figure 2d, bottom panel). The intraclass correlation coefficient across all cohorts was moderate (.218), demonstrating that variation within the dataset was significantly driven by natural variation due to random X-chromosome inactivation, over animal-to-animal variability (See Materials and Methods). Together, these results suggest that MeCp2 expression in non-PV cells affects PNN expression associated with PV cells in an age-dependent manner.

### 3.3 | Adolescent Het perform a pup retrieval task with comparable efficiency to age-matched WT

Adult Het display sensory deficits while performing the complex social behavior of alloparental care, as measured by the pup retrieval task (Krishnan et al., 2017; Stevenson et al., 2021). Since adolescent Het expressed more MeCp2 in non-PV nuclei and expressed PNNs at WT levels, we predicted they would perform efficiently in the pup retrieval task (Figure 3). Adolescent Het and WT improved at pup retrieval across six consecutive days of testing as measured by time to retrieve scattered pups to the nest (latency index; Figure 3a,a') and number of errors (Figure 3b,b'). Additionally, adolescent Het retrieved pups with a similar efficiency as adolescent WT females. Adult Het are comparably inefficient at this same task (Stevenson et al., 2021). Cumulatively, these results suggest that the increased MeCp2 levels within non-PV nuclei, as well as typical PNN expression associated

TABLE 1 Definitions of parameters used to score tactile behaviors.

Whisker	Mouse within 3 cm of the object (due to typical whisker length) and actively exploring around texture/object
Whole-body climbing	All four paws are off the ground and animal is climbing the texture/object
Forepaw	Touching the object/texture with forepaws while the hind limbs are contacting the ground

TABLE 2 Number of animals included and excluded from analysis.

		Included in analysis		Excluded from analysis	
		WT	Het	WT	Het
12 WO	Texture discrimination day 1 (trials 1 & 2)	8	7	1	1
	Texture discrimination day 2 (trials 3 & 4)	9	8	1	0
	Novel object recognition (trials 1 & 2)	9	8	0	0
6 WO	Texture discrimination day 1 (trials 1 & 2)	8	8	1	2
	Texture discrimination day 2 (trials 3 & 4)	9	10	1	0
	Novel object recognition (trials 1 & 2)	9	10	0	0

Note: Animals were excluded from analysis due to technical difficulties with video lagging/skipping or no interactions with textures.

with PV+ interneurons, provide a molecular mechanism for compensatory behavioral adaptations during adolescence, which are lost in older females, resulting in a regressive behavioral phenotype.


### 3.4 | Het display regression in whisker tactile sensory behavior between adolescence and adulthood

Pup retrieval involves initial perception of sensory cues, multisensory integration, and motor coordination involving multiple neural circuits. Here, we chose to focus specifically on the single sensory perception modality of whisker tactile perception due to its connection to S1BF. We performed standard texture discrimination and object recognition tasks on adolescent and adult WT and Het and analyzed their behavior frame by frame using DataVyu software (Figure 4).


Standard representation of percentage of time spent with different textures did not significantly change in adolescent females

during texture investigation (WT:  $50.14\% \pm 4.344$ ,  $n=9$ ; Het:  $55.60\% \pm 4.116$ ,  $n=10$ ; one-sample *t*-test with 50% chance) or during object investigation (WT:  $47.74\% \pm 4.263$ ,  $n=9$ ; Het:  $50.33\% \pm 3.572$ ,  $n=10$ ; one-sample *t*-test with 50% chance). Percentage of time spent with textures did not change in adult WT or Het female mice (WT:  $57.08\% \pm 3.356$ ,  $n=9$ ; Het:  $51.59\% \pm 3.531$ ,  $n=8$ ; one-sample *t*-test with 50% chance) and during object investigation (WT:  $51.67\% \pm 4.477$ ,  $n=9$ ; Het:  $57.26\% \pm 4.511$ ,  $n=8$ ; one-sample *t*-test with 50% chance). Thus, we refrain from drawing conclusions about preferences of the female WT and Het mice or their memory capabilities here.


We chose to perform a systematic analysis of behavioral videos, as mice interact with textures and objects with high speed and rarely linger during investigations. We found that both adolescent WT and Het females predominantly investigated textures with whiskers and interacted with similar number of contacts (Figure 5a,b). However, though the total interaction time with textures did not differ significantly between adolescent WT and Het (Figure 5c), the duration per whisker interaction was significantly higher in Het (Figure 5d).



**FIGURE 3** Adolescent Het perform pup retrieval comparable to WT. (a–a') Mean latency index of pup retrieval behavior for each testing day (a) and averaged per animal across all testing days (a') showed significant improved retrieval across testing days for both WT and Het, and no significant difference in average latency between genotypes. (b–b') Mean number of errors during pup retrieval for each testing day (b) and averaged per animal across all testing days (b') showed same result as latency index. (All panels) mean  $\pm$  SEM are shown. (a, b) Kruskal–Wallis followed by uncorrected Dunn's test; \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ , \*\*\*\* $p < .0001$ . Color \* = within genotype, significance between postnatal day 0 (P0) and P-day. (a', b') Mann–Whitney test: ns, not significant.



**FIGURE 4** Tactile behavior schema. Behavior tests begin with 2 days of habituation to introduce the mice to the behavioral box setup. On each day of texture discrimination (days 3 and 4), individual mouse undergoes a 10-min familiarization trial followed by a 10-min testing trial, with varied sandpaper grits. On day 5, objects are used instead of textures for the novel object recognition task. See Materials and Methods for detail.




**FIGURE 5** Adolescent Het have subtle whisking deficits during texture interactions. (a) Examples of frame-by-frame texture interactions across four 10-min trials of texture exposure for adolescent WT (top panel) and Het (bottom panel), represented as sequences. The X-axis represents the time (s). Whisker interactions (purple) are plotted as blocks that span the duration of the interaction. Each row depicts fleeting whisker interactions of a mouse during individual trials. (b–d) Statistical analysis of the interaction quantifications revealed that on average, WT and Het had similar total number of contacts across two to four trials (b) and spent similar averaged total time investigating textures across two to four trials (c). However, Het spent significantly longer during each whisker interaction than WT (d). Superplots data visualization shows individual cohorts by same color circles, represent the mean of two to four trials ( $n=9$  WT,  $n=10$  Het) and violin plots show median and distribution of all interactions. Mann–Whitney test: ns, not significant; \*\*\*\* $p < .0001$ .


We performed similar analyses with novel object recognition to determine if adolescent Het are deficient in other forms of tactile sensory-dependent tasks. Adolescent WT and Het used their whiskers, forepaws, and whole-body climbing to investigate objects (Figure 6a). Interestingly, adolescent Het made significantly fewer whisker contacts (Figure 6b) and spent similar total time investigating objects (Figure 6e) compared to age-matched WT. There was no significant difference in the number of contacts or time spent investigating objects using forepaws (Figure 6c,f) and whole-body climbing (Figure 6d,g). Similar to texture investigation, adolescent Het interacted significantly longer per interaction using whiskers (Figure 6h), but not forepaws or whole-body climbing (Figure 6i,j), compared to WT. Together, the increased duration of interactions when interacting with textures and objects suggests the beginnings of an atypical whisker perception in adolescent Het.

On the contrary, adult Het displayed significant whisker hyposensitivity in texture discrimination compared to adult WT (Figure 7). Examples of frame-by-frame sequence analysis showed

that non-whisker interactions (forepaw and whole-body climbing) were interspersed with whisker interactions in adult Het, while WT used only whiskers (Figure 7a). Compared to WT, adult Het used whisker to make significantly more contacts, spent more time investigating textures, and had longer durations per whisker interaction with textures (Figure 7b,e,h). Adult Het also used forepaws and whole-body climbing to make significantly more contacts, spent more time investigating, and had longer time per interaction with textures than adult WT (Figure 7c,d,f,g,i,j, respectively). These findings are similar to previous studies in whiskerless and barrel-less mice (Guic-Robles et al., 1992; Guic-Robles et al., 1989) and suggest adult Het are seeking compensatory tactile sensation due to deficits in whisker sensation (hyposensitivity). Furthermore, this systematic analysis captured the individual variation expected in adult Het phenotypes. Adult Het 3 and 5 (Figure 7b–d,h–j, light blue and yellow colors, respectively) were similar to adult WT in number and duration of contacts with whiskers, and no interactions with forepaw or whole body, suggesting a likely typical functionality



**FIGURE 6** Adolescent Het exhibit subtle whisking deficits during object interactions. (a) Examples of frame-by-frame texture interactions across two 10-min trials of object exposure for adolescent WT (top panel) and Het (bottom panel) color coded by types of interaction (whisker, forepaw, whole-body climbing), represented as sequences. The X-axis represents the time (s), and interactions are plotted as blocks that span the duration of the interaction. Each row depicts fleeting whisker (purple), forepaw (yellow), or whole-body climbing (teal) interactions of the mice during individual trials. (b–d) Statistical analysis of the interaction quantifications revealed Het made significantly more contacts using whiskers (b), but not forepaws (c) or whole-body climbing (d) compared to WT. (e–g) There were no statistical differences between WT and Het for the time spent with objects using whiskers (e), forepaws (f), or whole-body climbing (g). (h–j) Het spent significantly more time than WT during each whisker interaction (h), but not with forepaw (i) or whole-body climbing (j). (b–j) Superplots data visualization shows individual cohorts by same color circles, represent the mean of two trials ( $n=9$  WT,  $n=10$  Het), and violin plots show median and distribution of all interactions. Mann–Whitney test: ns, not significant, \*\* $p < .01$ , \*\*\*\* $p < .0001$ .



**FIGURE 7** Adult Het exhibit hyposensitivity and compensatory behaviors during texture interactions. (a) Examples of frame-by-frame texture interactions across four 10-min trials of texture exposure for adult WT (top panel) and Het (bottom panel) color coded by types of interaction (whisker, forepaw, whole-body climbing), represented as sequences. The X-axis represents the time (s), and interactions are plotted as blocks that span the duration of the interaction. Each row depicts fleeting whisker (purple), forepaw (yellow), or whole-body climbing (teal) interactions of the mice during individual trials. (b–j) Statistical analysis of the interaction quantifications revealed adult Het made significantly more contacts (b–d), spent significantly more total time (e–g) and time per interaction (h–j) with textures, with all interaction types (b–j), compared to adult WT. Superplots data visualization shows individual cohorts by same color circles, represent the mean of two to four trials ( $n=9$  WT,  $n=8$  Het) and violin plots show median and distribution of all interactions. Mann–Whitney test: \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ , \*\*\*\* $p < .0001$ .

within S1BF of the respective animals. Adult Het 6 and 7 (dark blue and deep orange, respectively) performed forepaw interactions, but not whole-body climbing (Figure 7c,d,f,g,i,j), suggesting

minor sensory processing deficits in S1BF that is compensated for by appropriate tactile perception within the forelimb region of the primary somatosensory cortex. The other five adult Het were most

different from adult WT in their texture interactions and are likely to have profound deficits in neural processing in S1BF and neighboring SS1 subregions.

Surprisingly, adult Het had milder phenotypes while interacting with objects (Figure 8). Unlike texture investigations in adults, both adult WT and Het also used forepaws and whole-body climbing to interact with objects (Figure 8a). We did not observe significant differences in both the total number of contacts with objects and the total time spent investigating objects between adult WT and Het during whisker, forepaw, or whole-body climbing interactions (Figure 8b–g). Adult Het spent significantly longer time per interaction with objects using whiskers (Figure 8h), but not forepaws or whole-body climbing (Figure 8i,j). Taken together, these results point to the beginnings of a regressive phenotype associated with specific tactile sensory perception in 6-week-old female Het that progresses to a severe, context-dependent tactile hyposensitivity by 12 weeks of age.

### 3.5 | Context-specific developmental progression in WT and regression in Het

To develop a baseline in tactile performance of females in both adolescence and adulthood, we compared texture- and object-associated phenotypes within genotype and across age. Compared to adolescent WT, we found that adult WT spent significantly less time with textures (Figure 9a,b), while the number of contacts remained similar across ages (Figure 9c). Adult Het spent significantly more time with textures and more numbers of contacts (Figure 9d,f) compared to adolescent Het, while the duration per contact remained similar (Figure 9e). Together, these results suggest a progressive perception via whisker tactile interactions over age in WT, while Het display regressive inefficiencies by these same parameters. Interestingly, neither WT nor Het showed any significant age-specific changes in interactions with objects (Figure 10a–f), suggesting that female mice may use different strategies while interacting with finer textures and objects.

## 4 | DISCUSSION

Disrupted sensory processing is a contributing factor to many of the behavioral phenotypes in neurodevelopmental disorders (Rais et al., 2018; Schaffler et al., 2022). Here, we use established models of whisker tactile sensory perception to discover the impact of mosaic MeCp2 expression in identified cell types across developmental stages in a female mouse model for Rett syndrome (Figure 11). These results set the stage to explore the mechanistic links between age-dependent MeCp2 expression levels and neural circuits involved in sensory perception during simple and complex behavioral tasks within Het. Two major observations are of importance to the RTT field: (1) wild-type MeCp-2 protein expression is regulated in an age- and cell-type specific manner in Het, providing an unexplored angle for therapeutic intervention in the Het brain via dynamic protein regulation, and (2) female Het mice have typical functioning capacity which is

lost with age, suggesting that the Het brain is capable of cellular and behavioral plasticity, even past early infancy. Deep phenotyping at cellular and behavioral levels is necessary in this important preclinical model of Rett syndrome to aid in better therapeutic treatments.


### 4.1 | Aberrant homeostasis of MeCp2 in late adolescence as a catalyst for the transition from pre-symptomatic to symptomatic RTT female mice

One of the most intriguing results here is the observed differences in MeCp2 expression within nuclei of Layer IV S1BF non-PV cells in adolescent Het compared to age-matched WT. The observation of cell-type-specific and age-dependent MeCp2 expression dynamics within Het provides new avenues for consideration in RTT pathogenesis. To date, Rett syndrome is thought to be caused by the lack of functional MeCp2 in roughly half of all cells. We speculate that the transient fluctuation in MeCp2 levels within the ~50% of the cells expressing the wild-type protein is an additional direction to consider. The precocious increase in MeCp2 within non-PV nuclei of Het may be a cell-autonomous mode of compensation for the lack of MeCp-2 protein in surrounding cells. Over time, however, the ability to further increase MeCp2 expression levels in non-PV nuclei, as observed in all WT cells as well as Het PV cells, is lost.


Thus, although the genotype and the cellular status of the Het remain the same throughout life, their MeCp2 expression levels are changing in a cell-type specific manner, suggesting built-in homeostatic mechanisms to regulate MeCp2 expression. This could be a basis for normal development in infancy for girls diagnosed with RTT, and then regression at different times throughout life (Neul, 2019). It is equally intriguing that MeCp2 expression in PV neurons, with higher basal levels than in non-PV cells, exhibits typical increases similar to WT females across age. Yet, the underlying mechanisms regulating MeCp2 expression levels within specific cell types across age remain unclear. Sorting specific cell types and measuring MeCp-2 protein content, with regional and age specificities, are necessary to establish an extensive expression baseline in Het, as well as to determine how MeCp2 expression changes with experience within context-specific circuitry.

In female Het mice, the precocious, cell-type specific increases in MeCp2 expression coincide with pubertal changes. In female WT mice, onset of puberty is accelerated by leptin (Ahima et al., 1997; Chehab et al., 1997) and ovarian hormones that regulate the organization and maturation of cortices by regulating inhibition (Clemens et al., 2019; Piekarski et al., 2017). This line of research is important as girls with RTT enter puberty early and reach menarche later, and body mass index affects pubertal timing (Killian et al., 2014). If Het mice also undergo pubertal deviations from their WT littermates, atypical neuroendocrine signaling would affect Het females as they transition into adulthood, in terms of both steroid-dependent organization of circuitry and MeCp2 expression.

MeCp2 expression changes in appropriate primary sensory cortices in early developmental critical periods (He et al., 2014;



**FIGURE 8** Adult Het display mild whisker deficit while interacting with objects. (a) Examples of frame-by-frame texture interactions across two 10-min trials of object exposure for adult WT (top panel) and Het (bottom panel) color coded by types of interaction (whisker, forepaw, whole-body climbing), represented as sequences. The X-axis represents the time (s), and interactions are plotted as blocks that span the duration of the interaction. Each row depicts fleeting whisker (purple), forepaw (yellow), or whole-body climbing (teal) interactions of the mice during individual trials. (b–j) Statistical analysis of the interaction quantifications revealed adult Het interacted with objects using whisker, forepaw, and whole-body climbing, similar to WT, in the number of contacts (b–d), total time with objects (e–g), and time per interactions (with Het having a slight but significant increase with whisker) (h–j). Superplots show animals color coded by circles, represent the mean of two trials ( $n=9$  WT,  $n=8$  Het). Superimposition of violin plots show median and distribution of all interactions. Mann–Whitney test: \*\* $p < .01$ ; ns, not significant.




**FIGURE 9** Differential and age-specific changes in WT and Het during texture investigation. (a–c) Using whiskers, adult WT interacted significantly shorter time (a: total time, b: time per interaction) with textures than adolescent WT, without changes in number of contacts made (c). (d–f) However, adults Het used whiskers to spent significantly longer time (d) and made more contacts (f) with textures than adolescent Het, without changes in time spent per interaction (e). (a–f) Each circle represents the mean of two to four trials per animal (adolescent:  $n=9$  WT,  $n=10$  Het; adult:  $n=9$  WT,  $n=8$  Het). Superimposition of violin plots shows median and distribution of all interactions. Mann–Whitney test:  $*p < .05$ ,  $****p < .0001$ ; ns, not significant.

Krishnan et al., 2015; Skene et al., 2010; Yagasaki et al., 2018) and in response to different stimuli (Martínez de Paz et al., 2019; Yagasaki et al., 2018). MeCp2 expression in adolescence and adulthood is thought to be stable. However, systematic studies with cell-type specific fluorescent-assisted cell sorting over brain regions, ages, and experimental conditions are required to determine the extent of the dynamic range of MeCp2 expression. Does the expression level of MeCp2 matter for its nuclear functions of binding methylated DNA and regulating gene expression? Increasing MeCp2 expression in transgenic overexpression rodent lines (Collins et al., 2004) or deleting MeCp2 in specific cell types also largely results in Rett syndrome-like phenotypes at behavioral and cellular levels (Bhattacharjee et al., 2017; Chapleau et al., 2009; Goffin et al., 2012; He et al., 2014; Ito-Ishida et al., 2015; Krishnan et al., 2017; Meng et al., 2016; Mossner et al., 2020). At the molecular level, cells that do not express MeCp2 could fail to maintain precise and stable gene expression programs over time, that could also contribute to the regressive phenotypes (Johnson et al., 2017; Linhoff et al., 2015; Renthal et al., 2018; Zhao et al., 2013). In general, how MeCp2 expression in neighboring cells of Het is affected is unknown in these manipulations. Moreover, considering the possibility of compensatory changes in MeCp2 expression, it is critical to determine the impact of these manipulations on MeCp2 expression in unmanipulated,

non-target cells in an age-specific manner. Although these MeCp2 expression changes are identified using well-established immunostaining and confocal microscopy techniques, to obtain single cell protein expression with spatial resolution, further validations using other proteomics approaches would be helpful to validate these changes. Developing these methods would be very useful, as we speculate such compensatory mechanisms at the protein level play crucial roles in other X-linked neurodevelopmental disorders (Brand et al., 2021; Santos-Rebouças et al., 2020) and metabolic disorders (Juchniewicz et al., 2021).

#### 4.2 | Perineuronal nets and PV-pyramidal networks in *Mecp2*-deficient mouse models

Germline deletion of *Mecp2* from all cell types as well as conditional deletion of *Mecp2* from both PV neurons and forebrain pyramidal neurons result in myriad of cellular and behavioral phenotypes, particularly in male mice (Chao et al., 2010; Gemelli et al., 2006; Ito-Ishida et al., 2015; Krishnan et al., 2015; Morello et al., 2018; Mossner et al., 2020; Patrizi et al., 2020). An altered PV-network configuration interferes with experience-dependent plasticity mechanism in the brain and is associated with defective learning




**FIGURE 10** No differential or age-specific changes in WT and Het during object interaction. (a–c) Both adolescent and adult WT used whiskers similar to interact with objects, as determined by total time spent (a), time per interaction (b), and number of contacts (c). (d–f) Comparably, adolescent and adult Het used whiskers similar to interact with objects (d: total time; e: time per interaction; f: number of contacts). Each circle represents the mean of two trials per animal (adolescent:  $n=9$  WT,  $n=10$  Het; adult:  $n=9$  WT,  $n=8$  Het). Superimposition of violin plots show median and distribution of interactions. Mann–Whitney test: ns, not significant.

and plasticity (Donato et al., 2013; Krishnan et al., 2015; Patrizi et al., 2020). Adult Het mice have atypical increased expression of PNNs in the auditory cortex (Krishnan et al., 2017) and specific subregions of the primary somatosensory cortex (Lau, Layo, et al., 2020). Typically, pup vocalizations trigger suppression of PV+ GABAergic neuronal responses and a concomitant disinhibition in deep-layer pyramidal neurons in the auditory cortex of adult WT surrogates (Lau, Krishnan, et al., 2020). In adult Het surrogates, the increased perineuronal net expression interferes with the synaptic plasticity of PV+ GABAergic neurons, resulting in a lack of disinhibition of pyramidal neurons (Lau, Krishnan, et al., 2020) and coinciding with poor pup retrieval performance (Krishnan et al., 2017). It is currently unknown if similar mechanisms also occur in the primary somatosensory cortex, and if so, whether they occur in an age-dependent manner, which would ultimately lead to atypical tactile processing in Het. Studies are underway to determine how PNN expression in S1BF contributes to tactile sensory perception.

### 4.3 | Systematic analysis of behavioral assays to study tactile sensory perception in female mice

Traditionally, texture discrimination and object recognition assays are reported as “percent novelty recognition” or a “discrimination

index” calculated as total time spent investigating the novel texture/object divided by the total time spent investigating both textures/objects (Antunes & Biala, 2012; Ennaceur et al., 2005; Ennaceur & Delacour, 1988). This analysis normalizes the time spent investigating across individual animals to control variation in exploration time. Based on these metrics, a mixed result of tactile hyper- and hyposensitivity has been previously reported in models of MeCp2 deficiency (Chao et al., 2010; Chen et al., 2001; Gemelli et al., 2006; Ito-Ishida et al., 2015; Orefice et al., 2016). It is hypothesized that mice will spend more time with novel texture/object due to their innate nature of curiosity and their ability to remember old or novel objects/textures from such interactions. However, a growing body of literature shows that freely moving rodents do not have a preference for novelty (Antunes & Biala, 2012; Cohen & Stackman Jr., 2015; Ennaceur, 2010). Mice and rats show preference for familiar objects rather than novel objects (Ennaceur, 2010). Adult C57BL/6 male mice show novelty object preference, whereas C57BL/6 adult females show no preference between a familiar and novel object (Frick & Gresack, 2003), consistent with our findings. Thus, to reconcile these differences in the literature and establish first-order metrics for analysis, we performed a systematic, frame-by-frame analysis of interactions with textures and objects for adolescent and adult female C57BL/6 mice. By analyzing the number and time



**FIGURE 11** Summary diagram: MeCp2 and PNN expression coincide with age-dependent sensory deficits in Het. Wild-type MeCp2 expression is increased in genotypic adolescent *Mecp2*-heterozygous female mice, which coincides with typical expression of perineuronal nets in barrel field of the primary somatosensory cortex, mild tactile sensory perception, and efficient multisensory processing during the pup retrieval task. In adulthood, wild-type MeCp2 expression in Het is similar to control WT brains, which coincides with atypical increases in perineuronal nets, increased whisker interactions indicative of hyposensitivity, and aberrant multisensory processing leading to inefficient pup retrieval.

of interactions, we found that adult WT females spent less time interacting with textures compared to adolescent WT females. Adult Het females interacted more overall and longer than adolescent Het females, suggesting a regressive tactile hyposensitivity phenotype in adult Het.

#### 4.4 | Regression in RTT models

Previous work using a female rat RTT model has reported a regression in a learned forepaw skill over the course of weeks (Veeraragavan et al., 2016). Regressive breathing abnormalities have also been shown using a conditional MeCp2 knockout male mouse model (Huang et al., 2016). Previously, we have proposed that adult female Het show regression in maintaining pup retrieval efficiency over few days (Stevenson et al., 2021). While these studies report regression in parasympathetic and motor function, our work here is the first to report a regressive hyposensitivity phenotype in tactile perception between adolescence and adulthood in a female mouse model for RTT. An alternative explanation for increases in tactile interactions from adolescence to adulthood in Het is the emergence of repetitive obsessive-compulsive disorder-like phenotypes over time, as observed in mouse models of this disorder (Ahmari, 2016; Alonso et al., 2015; Mitra & Bult-Ito, 2021). Adult Het perform repetitive

jumping during early days of pup retrieval which could be indicative of novelty- or stress-induced stereotypies in a subpopulation of adult Het (Stevenson et al., 2021). It is currently unclear if the increased frequency and longer durations of interactions in adult Het are a form of repetitive obsessive-compulsive behavior. Clozapine injection has been shown to exacerbate such repetitive phenotypes (Kang et al., 2020) and can be utilized to test this hypothesis in the future.

#### 4.5 | Rehabilitation strategies

Early interventions aimed at increasing neuronal activity mitigate disease onset in both male and female mouse models for Rett syndrome. Early administration of Ampakine CX546, a positive modulator of AMPA receptors, from postnatal days 3–9 in male *Mecp2*-null mice enhances neuronal activity, prolongs lifespan, delays disease progression, and rescues both motor abilities and spatial memory (Scaramuzza et al., 2021). In Het females, repetitive motor training enhances neuronal activity of task-specific neurons and improves both spatial memory and motor function well into adulthood (Achilly et al., 2021). Adolescent and adult Het improve in the pup retrieval task with repeated trials over days, in contrast to nonconsecutive days of training (Krishnan et al., 2017;

Stevenson et al., 2021). Early and repetitive trainings delay the onset of disease phenotypes and improve the latency to survival in Het, supporting the plastic capacity of the adolescent Het brain (Achilly et al., 2021). We speculate that wild-type MeCp2 expression in Het is affected by these trainings, especially at earlier ages, resulting in activation of compensatory pathways which lead to appropriate behavioral adaptations. Further work is needed to identify therapeutic and rehabilitative strategies that target MeCp2 maintenance within the female Het brain for mitigating disease progression throughout life.

#### DECLARATION OF TRANSPARENCY

The authors, reviewers and editors affirm that in accordance to the policies set by the *Journal of Neuroscience Research*, this manuscript presents an accurate and transparent account of the study being reported and that all critical details describing the methods and results are present.

#### AUTHOR CONTRIBUTIONS

All authors had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. *Conceptualization*, K.K. and D.L.C.; *Methodology*, K.K., B.Y.B.L., D.L.C., M.M., and L.D.; *Software*, M.M., T.H., and A.W.; *Validation*, B.Y.B.L.; *Investigation*, D.L.C., D.W.S., L.D., M.M., T.R.S., A.M., S.P., and B.Y.B.L.; *Formal Analysis*, L.D., M.M., A.W., and T.H.; *Resources*, K.K.; *Data Curation*, D.L.C., M.M., L.D., A.W., and T.H.; *Writing – Original Draft*, M.M.; *Writing – Review & Editing*, K.K., M.M., L.D., D.L.C., A.W., T.H., and B.Y.B.L.; *Visualization*, M.M., L.D., A.W., and T.H.; *Supervision*, K.K.; *Project Administration*, K.K.; *Funding Acquisition*, K.K., B.Y.B.L., M.M., D.L.C., L.D., and T.H.

#### ACKNOWLEDGMENTS

We would like to thank undergraduates Beyza Kartal, Benjamin Alireza Bridges, Sami Issac, Shahin Ahmadi, Itzanami Sotelo Hernandez, Miranda Blevins, and Deema Mansour for technical help, and the 2020 and 2021 BCMB 420 Image analysis class for help in mapping and analyzing high-intensity PNN data. We would like to thank the lab of Dr. Maitreyi Das and her graduate students for training and assistance using the confocal microscope. This work was supported by BCMB Chancellor's fellowships (MM and DLC), the 2021 Scholarly and Research Incentive Funds (SARIF) through the Office of Research, Innovation, & Economic Development (ORIED) at UTK (MM), BCMB James and Dora Wright Fellowship (MM), W. Sherman Kouns Excellence in Teaching award (MM), National Science Foundation Graduate Research Fellowship Program (GRFP) (LD), postdoctoral fellowship award from [Rettsyndrome.org](https://www.rett.org/) (BYBL), the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R01GM140462 (TH), startup funds from the University of Tennessee–Knoxville (KK), and by the National Institute of Mental Health of the National Institutes of Health under Award Number R15MH124042 (KK).

#### CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

#### PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1002/jnr.25190>.

#### DATA AVAILABILITY STATEMENT

Data from this study will be deposited on Zenodo. The reserved DOI is <https://doi.org/10.5281/zenodo.7576575>, which will be activated after manuscript acceptance.

#### ORCID

Michael Mykins  <https://orcid.org/0000-0003-3467-5308>  
 Dana Layo-Carris  <https://orcid.org/0000-0001-7009-0652>  
 Logan Reid Dunn  <https://orcid.org/0000-0002-2147-5502>  
 Andrew Willems  <https://orcid.org/0000-0002-7927-0745>  
 Billy You Bun Lau  <https://orcid.org/0000-0002-2380-5341>  
 Tian Hong  <https://orcid.org/0000-0002-8212-7050>  
 Keerthi Krishnan  <https://orcid.org/0000-0002-0858-4624>

#### REFERENCES

- Achilly, N. P., Wang, W., & Zoghbi, H. Y. (2021). Presymptomatic training mitigates functional deficits in a mouse model of Rett syndrome. *Nature*, 592(7855), 596–600. <https://doi.org/10.1038/s41586-021-03369-7>
- Ahima, R. S., Dushay, J., Flier, S. N., Prabakaran, D., & Flier, J. S. (1997). Leptin accelerates the onset of puberty in normal female mice. *Journal of Clinical Investigation*, 99(3), 391–395. <https://doi.org/10.1172/JCI119172>
- Ahmari, S. E. (2016). Using mice to model obsessive compulsive disorder: From genes to circuits. *Neuroscience*, 321, 121–137. <https://doi.org/10.1016/j.neuroscience.2015.11.009>
- Alonso, P., López Solà, C., Real, E., Segalàs, C., & Menchon, J. M. (2015). Animal models of obsessive-compulsive disorder: Utility and limitations. *Neuropsychiatric Disease and Treatment*, 11, 1939–1955. <https://doi.org/10.2147/NDT.S62785>
- Alsina-Llanes, M., De Brun, V., & Olazábal, D. E. (2015). Development and expression of maternal behavior in naïve female C57BL/6 mice: Ontogeny of maternal behavior in mice. *Developmental Psychobiology*, 57(2), 189–200. <https://doi.org/10.1002/dev.21276>
- Amir, R. E., Van den Veyver, I. B., Wan, M., Tran, C. Q., Francke, U., & Zoghbi, H. Y. (1999). Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. *Nature Genetics*, 23(2), 185–188. <https://doi.org/10.1038/13810>
- Antunes, M., & Biala, G. (2012). The novel object recognition memory: Neurobiology, test procedure, and its modifications. *Cognitive Processing*, 13(2), 93–110. <https://doi.org/10.1007/s10339-011-0430-z>
- Beach, F. A., & Jaynes, J. (1956). Studies of maternal retrieving in rats. III. Sensory cues involved in the lactating female's response to her young. *Behaviour*, 10, 104–124.
- Bebbington, A., Anderson, A., Ravine, D., Fyfe, S., Pineda, M., de Klerk, N., Ben-Zeev, B., Yatawara, N., Percy, A., Kaufmann, W. E., & Leonard, H. (2008). Investigating genotype-phenotype relationships in Rett syndrome using an international data set. *Neurology*, 70(11), 868–875. <https://doi.org/10.1212/01.wnl.0000304752.50773.ec>
- Bhattacharjee, A., Mu, Y., Winter, M. K., Knapp, J. R., Eggmann, L. S., Gunewardena, S. S., Kobayashi, K., Kato, S., Krizsan-Agbas, D.,

- & Smith, P. G. (2017). Neuronal cytoskeletal gene dysregulation and mechanical hypersensitivity in a rat model of Rett syndrome. *Proceedings of the National Academy of Sciences of the United States of America*, 114(33), E6952–E6961. <https://doi.org/10.1073/pnas.1618210114>
- Bibbings, K., Harding, P. J., Loram, I. D., Combes, N., & Hodson-Tole, E. F. (2019). Foreground detection analysis of ultrasound image sequences identifies markers of motor neurone disease across diagnostically relevant skeletal muscles. *Ultrasound in Medicine & Biology*, 45(5), 1164–1175. <https://doi.org/10.1016/j.ultrasmedbio.2019.01.018>
- Bignami, A., Asher, R., & Perides, G. (1992). Co-localization of hyaluronic acid and chondroitin sulfate proteoglycan in rat cerebral cortex. *Brain Research*, 579(1), 173–177.
- Boxer, L. D., Renthal, W., Greben, A. W., Whitwam, T., Silberfeld, A., Stroud, H., Li, E., Yang, M. G., Kinde, B., Griffith, E. C., Bonev, B., & Greenberg, M. E. (2020). MeCP2 represses the rate of transcriptional initiation of highly methylated long genes. *Molecular Cell*, 77(2), 294–309.e9. <https://doi.org/10.1016/j.molcel.2019.10.032>
- Brand, B. A., Blesson, A. E., & Smith-Hicks, C. L. (2021). The impact of X-Chromosome inactivation on phenotypic expression of X-Linked neurodevelopmental disorders. *Brain Sciences*, 11(7), 904. <https://doi.org/10.3390/brainsci11070904>
- Braunschweig, D., Simcox, T., Samaco, R. C., & LaSalle, J. M. (2004). X-Chromosome inactivation ratios affect wild-type MeCP2 expression within mosaic Rett syndrome and *Mecp2*<sup>-/+</sup> mouse brain. *Human Molecular Genetics*, 13(12), 1275–1286. <https://doi.org/10.1093/hmg/ddh142>
- Buchanan, C. B., Stallworth, J. L., Scott, A. E., Glaze, D. G., Lane, J. B., Skinner, S. A., Tierney, A. E., Percy, A. K., Neul, J. L., & Kaufmann, W. E. (2019). Behavioral profiles in Rett syndrome: Data from the natural history study. *Brain and Development*, 41(2), 123–134. <https://doi.org/10.1016/j.braindev.2018.08.008>
- Carstens, K. E., Lustberg, D. J., Shaughnessy, E. K., McCann, K. E., Alexander, G. M., & Dudek, S. M. (2021). Perineuronal net degradation rescues CA2 plasticity in a mouse model of Rett syndrome. *Journal of Clinical Investigation*, 131(16), e137221. <https://doi.org/10.1172/JCI137221>
- Carstens, K. E., Phillips, M. L., Pozzo-Miller, L., Weinberg, R. J., & Dudek, S. M. (2016). Perineuronal nets suppress plasticity of excitatory synapses on CA2 pyramidal neurons. *Journal of Neuroscience*, 36(23), 6312–6320. <https://doi.org/10.1523/JNEUROSCI.0245-16.2016>
- Carulli, D., Rhodes, K. E., Brown, D. J., Bonnert, T. P., Pollack, S. J., Oliver, K., Strata, P., & Fawcett, J. W. (2006). Composition of perineuronal nets in the adult rat cerebellum and the cellular origin of their components. *Journal of Comparative Neurology*, 494(4), 559–577. <https://doi.org/10.1002/cne.20822>
- Carulli, D., Rhodes, K. E., & Fawcett, J. W. (2007). Upregulation of aggrecan, link protein 1, and hyaluronan synthases during formation of perineuronal nets in the rat cerebellum. *Journal of Comparative Neurology*, 501(1), 83–94. <https://doi.org/10.1002/cne.21231>
- Chahrour, M., Jung, S. Y., Shaw, C., Zhou, X., Wong, S. T. C., Qin, J., & Zoghbi, H. Y. (2008). MeCP2, a key contributor to neurological disease, activates and represses transcription. *Science*, 320(5880), 1224–1229. <https://doi.org/10.1126/science.1153252>
- Champagne, F. A., & Curley, J. P. (2016). Plasticity of the maternal brain across the lifespan: Plasticity of the maternal brain across the lifespan. *New Directions for Child and Adolescent Development*, 2016(153), 9–21. <https://doi.org/10.1002/cad.20164>
- Champagne, F. A., Curly, J. P., Keverne, E. B., & Bateson, P. P. G. (2007). Natural variations in postpartum maternal care in inbred and outbred mice. *Physiology and Behavior*, 91, 325–334.
- Chao, H.-T., Chen, H., Samaco, R. C., Xue, M., Chahrour, M., Yoo, J., Neul, J. L., Gong, S., Lu, H.-C., Heintz, N., Ekker, M., Rubenstein, J. L. R., Noebels, J. L., Rosenmund, C., & Zoghbi, H. Y. (2010). Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature*, 468(7321), 263–269. <https://doi.org/10.1038/nature09582>
- Chapleau, C. A., Calfa, G. D., Lane, M. C., Albertson, A. J., Larimore, J. L., Kudo, S., Armstrong, D. L., Percy, A. K., & Pozzo-Miller, L. (2009). Dendritic spine pathologies in hippocampal pyramidal neurons from Rett syndrome brain and after expression of Rett-associated MECP2 mutations. *Neurobiology of Disease*, 35(2), 219–233. <https://doi.org/10.1016/j.nbd.2009.05.001>
- Charman, T., Cass, H., Owen, L., Wigram, T., Slonims, V., Weeks, L., Wisbeach, A., & Reilly, S. (2002). Regression in individuals with Rett syndrome. *Brain and Development*, 24(5), 281–283. [https://doi.org/10.1016/S0387-7604\(02\)00058-X](https://doi.org/10.1016/S0387-7604(02)00058-X)
- Chehab, F. F., Mounzih, K., Lu, R., & Lim, M. E. (1997). Early onset of reproductive function in normal female mice treated with leptin. *Science*, 275(5296), 88–90. <https://doi.org/10.1126/science.275.5296.88>
- Chen, L., Chen, K., Lavery, L. A., Baker, S. A., Shaw, C. A., Li, W., & Zoghbi, H. Y. (2015). MeCP2 binds to non-CG methylated DNA as neurons mature, influencing transcription and the timing of onset for Rett syndrome. *Proceedings of the National Academy of Sciences of the United States of America*, 112(17), 5509–5514. <https://doi.org/10.1073/pnas.1505909112>
- Chen, R. Z., Akbarian, S., Tudor, M., & Jaenisch, R. (2001). Deficiency of methyl-CpG binding protein-2 in CNS neurons results in a Rett-like phenotype in mice. *Nature Genetics*, 27(3), 327–331. <https://doi.org/10.1038/85906>
- Clemens, A. M., Lenschow, C., Beed, P., Li, L., Sammons, R., Naumann, R. K., Wang, H., Schmitz, D., & Brecht, M. (2019). Estrus-cycle regulation of cortical inhibition. *Current Biology*, 29(4), 605–615.e6. <https://doi.org/10.1016/j.cub.2019.01.045>
- Cohen, S. J., & Stackman, R. W., Jr. (2015). Assessing rodent hippocampal involvement in the novel object recognition task. A review. *Behavioural Brain Research*, 285, 105–117. <https://doi.org/10.1016/j.bbr.2014.08.002>
- Collins, A. L., Levenson, J. M., Vilaythong, A. P., Richman, R., Armstrong, D. L., Noebels, J. L., David Sweatt, J., & Zoghbi, H. Y. (2004). Mild overexpression of MeCP2 causes a progressive neurological disorder in mice. *Human Molecular Genetics*, 13(21), 2679–2689. <https://doi.org/10.1093/hmg/ddh282>
- de Vivo, L., Landi, S., Panniello, M., Baroncelli, L., Chierzi, S., Mariotti, L., Spolidoro, M., Pizzorusso, T., Maffei, L., & Ratto, G. M. (2013). Extracellular matrix inhibits structural and functional plasticity of dendritic spines in the adult visual cortex. *Nature Communications*, 4(1), 1484. <https://doi.org/10.1038/ncomms2491>
- Diamond, M. E., von Heimendahl, M., & Arabzadeh, E. (2008). Whisker-mediated texture discrimination. *PLoS Biology*, 6(8), e220. <https://doi.org/10.1371/journal.pbio.0060220>
- Djukic, A., & McDermott, M. V. (2012). Social preferences in Rett syndrome. *Pediatric Neurology*, 46(4), 240–242. <https://doi.org/10.1016/j.pediatrneurol.2012.01.011>
- Djukic, A., McDermott, M. V., Mavrommatis, K., & Martins, C. L. (2012). Rett syndrome: Basic features of visual processing—A pilot study of eye-tracking. *Pediatric Neurology*, 47(1), 25–29. <https://doi.org/10.1016/j.pediatrneurol.2012.04.009>
- Donato, F., Rompani, S. B., & Caroni, P. (2013). Parvalbumin-expressing basket-cell network plasticity induced by experience regulates adult learning. *Nature*, 504(7479), 272–276. <https://doi.org/10.1038/nature12866>
- Durand, S., Patrizi, A., Quast, K. B., Hachigian, L., Pavlyuk, R., Saxena, A., Carninci, P., Hensch, T. K., & Fagiolini, M. (2012). NMDA receptor regulation prevents regression of visual cortical function in the absence of *Mecp2*. *Neuron*, 76(6), 1078–1090. <https://doi.org/10.1016/j.neuron.2012.12.004>
- Einspieler, C., & Marschik, P. B. (2019). Regression in Rett syndrome: Developmental pathways to its onset. *Neuroscience & Biobehavioral Reviews*, 98, 320–332. <https://doi.org/10.1016/j.neubiorev.2019.01.028>

- Ennaceur, A. (2010). One-trial object recognition in rats and mice: Methodological and theoretical issues. *Behavioural Brain Research*, 215(2), 244–254. <https://doi.org/10.1016/j.bbr.2009.12.036>
- Ennaceur, A., & Delacour, J. (1988). A new one-trial test for neurobiological studies of memory in rats. 1: Behavioral data. *Behavioural Brain Research*, 31(1), 47–59. [https://doi.org/10.1016/0166-4328\(88\)90157-X](https://doi.org/10.1016/0166-4328(88)90157-X)
- Ennaceur, A., Michalikova, S., Bradford, A., & Ahmed, S. (2005). Detailed analysis of the behavior of Lister and Wistar rats in anxiety, object recognition and object location tasks. *Behavioural Brain Research*, 159(2), 247–266. <https://doi.org/10.1016/j.bbr.2004.11.006>
- Frick, K. M., & Gresack, J. E. (2003). Sex differences in the behavioral response to spatial and object novelty in adult C57BL/6 mice. *Behavioral Neuroscience*, 117(6), 1283–1291. <https://doi.org/10.1037/0735-7044.117.6.1283>
- Fuks, F., Hurd, P. J., Wolf, D., Nan, X., Bird, A. P., & Kouzarides, T. (2003). The methyl-CpG-binding protein MeCP2 links DNA methylation to histone methylation. *Journal of Biological Chemistry*, 278(6), 4035–4040. <https://doi.org/10.1074/jbc.M210256200>
- Gabel, H. W., Kinde, B., Stroud, H., Gilbert, C. S., Harmin, D. A., Kastan, N. R., Hemberg, M., Ebert, D. H., & Greenberg, M. E. (2015). Disruption of DNA-methylation-dependent long gene repression in Rett syndrome. *Nature*, 522(7554), 89–93. <https://doi.org/10.1038/nature14319>
- Gemelli, T., Berton, O., Nelson, E. D., Perrotti, L. I., Jaenisch, R., & Monteggia, L. M. (2006). Postnatal loss of methyl-CpG binding protein 2 in the forebrain is sufficient to mediate behavioral aspects of Rett syndrome in mice. *Biological Psychiatry*, 59(5), 468–476. <https://doi.org/10.1016/j.biopsych.2005.07.025>
- Goffin, D., Allen, M., Zhang, L., Amorim, M., Wang, I.-T. J., Reyes, A.-R. S., Mercado-Berton, A., Ong, C., Cohen, S., Hu, L., Blendy, J. A., Carlson, G. C., Siegel, S. J., Greenberg, M. E., & Zhou, Z. (2012). Rett syndrome mutation MeCP2 T158A disrupts DNA binding, protein stability and ERP responses. *Nature Neuroscience*, 15(2), 274–283. <https://doi.org/10.1038/nn.2997>
- Gogolla, N., Caroni, P., Lüthi, A., & Herry, C. (2009). Perineuronal nets protect fear memories from erasure. *Science*, 325(5945), 1258–1261. <https://doi.org/10.1126/science.1174146>
- Guic-Robles, E., Jenkins, W. M., & Bravo, H. (1992). Vibrissal roughness discrimination is barrelcortex-dependent. *Behavioural Brain Research*, 48(2), 145–152. [https://doi.org/10.1016/S0166-4328\(05\)80150-0](https://doi.org/10.1016/S0166-4328(05)80150-0)
- Guic-Robles, E., Valdivieso, C., & Guajardo, G. (1989). Rats can learn a roughness discrimination using only their vibrissal system. *Behavioural Brain Research*, 31(3), 285–289. [https://doi.org/10.1016/0166-4328\(89\)90011-9](https://doi.org/10.1016/0166-4328(89)90011-9)
- Guy, J., Hendrich, B., Holmes, M., Martin, J. E., & Bird, A. (2001). A mouse Mecp2-null mutation causes neurological symptoms that mimic Rett syndrome. *Nature Genetics*, 27(3), 322–326. <https://doi.org/10.1038/85899>
- Hagberg, B., Aicardi, J., Dias, K., & Ramos, O. (1983). A progressive syndrome of autism, dementia, ataxia, and loss of purposeful hand use in girls: Rett's syndrome: Report of 35 cases. *Annals of Neurology*, 14(4), 471–479. <https://doi.org/10.1002/ana.410140412>
- Hartig, W., Brauer, K., & Bruckner, G. (1992). Wisteria floribunda agglutinin-labelled nets surround parvalbumin-containing neurons. *Neuroreport*, 3(10), 869–872.
- He, L., Liu, N., Cheng, T., Chen, X., Li, Y., Shu, Y., Qiu, Z., & Zhang, X. (2014). Conditional deletion of Mecp2 in parvalbumin-expressing GABAergic cells results in the absence of critical period plasticity. *Nature Communications*, 5(1), 5036. <https://doi.org/10.1038/ncomms6036>
- Hemel, S. B. V. (1973). Pup retrieving as a reinforcer in nulliparous mice. *Journal of the Experimental Analysis of Behavior*, 19(2), 233–238. <https://doi.org/10.1901/jeab.1973.19-233>
- Hinde, R. A., & Davies, L. (1972). Removing infant rhesus from mother for 13 days compared with removing mother from infant. *Journal of Child Psychology and Psychiatry*, 13(4), 227–237. <https://doi.org/10.1111/j.1469-7610.1972.tb01149.x>
- Hou, X., Yoshioka, N., Tsukano, H., Sakai, A., Miyata, S., Watanabe, Y., Yanagawa, Y., Sakimura, K., Takeuchi, K., Kitagawa, H., Hensch, T. K., Shibuki, K., Igarashi, M., & Sugiyama, S. (2017). Chondroitin sulfate is required for onset and offset of critical period plasticity in visual cortex. *Scientific Reports*, 7(1), 12646. <https://doi.org/10.1038/s41598-017-04007-x>
- Huang, T.-W., Kochukov, M. Y., Ward, C. S., Merritt, J., Thomas, K., Nguyen, T., Arenkiel, B. R., & Neul, J. L. (2016). Progressive changes in a distributed neural circuit underlie breathing abnormalities in mice lacking MeCP2. *Journal of Neuroscience*, 36(20), 5572–5586. <https://doi.org/10.1523/JNEUROSCI.2330-15.2016>
- Ito-Ishida, A., Baker, S. A., Sillitoe, R. V., Sun, Y., Zhou, J., Ono, Y., Iwakiri, J., Yuzaki, M., & Zoghbi, H. Y. (2020). MeCP2 levels regulate the 3D structure of heterochromatic foci in mouse neurons. *Journal of Neuroscience*, 40(45), 8746–8766. <https://doi.org/10.1523/JNEUROSCI.1281-19.2020>
- Ito-Ishida, A., Ure, K., Chen, H., Swann, J. W., & Zoghbi, H. Y. (2015). Loss of MeCP2 in parvalbumin- and somatostatin-expressing neurons in mice leads to distinct Rett syndrome-like phenotypes. *Neuron*, 88(4), 651–658. <https://doi.org/10.1016/j.neuron.2015.10.029>
- Johnson, B. S., Zhao, Y.-T., Fasolino, M., Lamonica, J. M., Kim, Y. J., Georgakilas, G., Wood, K. H., Bu, D., Cui, Y., Goffin, D., Vahedi, G., Kim, T. H., & Zhou, Z. (2017). Biotin tagging of MeCP2 in mice reveals contextual insights into the Rett syndrome transcriptome. *Nature Medicine*, 23(10), 1203–1214. <https://doi.org/10.1038/nm.4406>
- Jones, P. L., Veenstra, G. J. C., Wade, P. A., Vermaak, D., Kass, S. U., Landsberger, N., Strouboulis, J., & Wolffe, A. P. (1998). Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nature Genetics*, 19(2), 187–191. <https://doi.org/10.1038/561>
- Juchniewicz, P., Piotrowska, E., Kloska, A., Podlacha, M., Mantej, J., Węgrzyn, G., Tukaj, S., & Jakóbkiewicz-Banecka, J. (2021). Dosage compensation in females with X-linked metabolic disorders. *International Journal of Molecular Sciences*, 22(9), 4514. <https://doi.org/10.3390/ijms22094514>
- Kang, S., Noh, H. J., Bae, S. H., Kim, Y.-S., Lew, H., Lim, J., Kim, S. J., Hong, K. S., Rah, J.-C., & Kim, C. H. (2020). Clozapine generates obsessive compulsive disorder-like behavior in mice. *Molecular Brain*, 13(1), 84. <https://doi.org/10.1186/s13041-020-00621-5>
- Kerr, A. (1995). Early clinical signs in the Rett disorder. *Neuropediatrics*, 26(2), 67–71. <https://doi.org/10.1055/s-2007-979725>
- Key, A. P., Jones, D., & Peters, S. (2019). Spoken word processing in Rett syndrome: Evidence from event-related potentials. *International Journal of Developmental Neuroscience*, 73(1), 26–31. <https://doi.org/10.1016/j.ijdevneu.2019.01.001>
- Killian, J. T., Lane, J. B., Cutter, G. R., Skinner, S. A., Kaufmann, W. E., Tarquinio, D. C., Glaze, D. G., Motil, K. J., Neul, J. L., & Percy, A. K. (2014). Pubertal development in Rett syndrome deviates from typical females. *Pediatric Neurology*, 51(6), 769–775. <https://doi.org/10.1016/j.pediatrneurol.2014.08.013>
- Kirby, R. S., Lane, J. B., Childers, J., Skinner, S. A., Annese, F., Barrish, J. O., Glaze, D. G., MacLeod, P., & Percy, A. K. (2010). Longevity in Rett syndrome: Analysis of the North American database. *Journal of Pediatrics*, 156(1), 135–138.e1. <https://doi.org/10.1016/j.jpeds.2009.07.015>
- Klose, R. J., Sarraf, S. A., Schmiedeberg, L., McDermott, S. M., Stancheva, I., & Bird, A. P. (2005). DNA binding selectivity of MeCP2 due to a requirement for A/T sequences adjacent to methyl-CpG. *Molecular Cell*, 19(5), 667–678. <https://doi.org/10.1016/j.molcel.2005.07.021>
- Koch, M., & Ehret, G. (1989). Estradiol and parental experience, but not prolactin are necessary for ultrasound recognition and

- pup-retrieving in the mouse. *Physiology & Behavior*, 45(4), 771–776. [https://doi.org/10.1016/0031-9384\(89\)90293-X](https://doi.org/10.1016/0031-9384(89)90293-X)
- Kosaka, T., & Heizmann, C. W. (1989). Selective staining of a population of parvalbumin-containing GABAergic neurons in the rat cerebral cortex by lectins with specific affinity for terminal N-acetylgalactosamine. *Brain Research*, 483(1), 158–163. [https://doi.org/10.1016/0006-8993\(89\)90048-6](https://doi.org/10.1016/0006-8993(89)90048-6)
- Krishnan, K., Lau, B. Y. B., Ewall, G., Huang, Z. J., & Shea, S. D. (2017). MECP2 regulates cortical plasticity underlying a learned behaviour in adult female mice. *Nature Communications*, 8(1), 14077. <https://doi.org/10.1038/ncomms14077>
- Krishnan, K., Wang, B.-S., Lu, J., Wang, L., Maffei, A., Cang, J., & Huang, Z. J. (2015). MeCP2 regulates the timing of critical period plasticity that shapes functional connectivity in primary visual cortex. *Proceedings of the National Academy of Sciences of the United States of America*, 112(34), E4782–E4791. <https://doi.org/10.1073/pnas.1506499112>
- Krususvee, V., Lyst, M. J., Taylor, C., Tarnauskaitė, Ž., Bird, A. P., & Cook, A. G. (2017). Structure of the MeCP2–TBLR1 complex reveals a molecular basis for Rett syndrome and related disorders. *Proceedings of the National Academy of Sciences of the United States of America*, 114(16), E3243–E3250. <https://doi.org/10.1073/pnas.1700731114>
- Kwok, J. C. F., Carulli, D., & Fawcett, J. W. (2010). In vitro modeling of perineuronal nets: Hyaluronan synthase and link protein are necessary for their formation and integrity: Hyaluronan synthase and link protein in PNNs. *Journal of Neurochemistry*, 114(5), 1447–1459. <https://doi.org/10.1111/j.1471-4159.2010.06878.x>
- Lau, B. Y. B., Krishnan, K., Huang, Z. J., & Shea, S. D. (2020). Maternal experience-dependent cortical plasticity in mice is circuit- and stimulus-specific and requires MECP2. *Journal of Neuroscience*, 40(7), 1514–1526. <https://doi.org/10.1523/JNEUROSCI.1964-19.2019>
- Lau, B. Y. B., Layo, D. E., Emery, B., Everett, M., Kumar, A., Stevenson, P., Reynolds, K. G., Cherosky, A., Bowyer, S.-A. H., Roth, S., Fisher, D. G., McCord, R. P., & Krishnan, K. (2020). Lateralized expression of cortical perineuronal nets during maternal experience is dependent on MECP2. *eNeuro*, 7(3), ENEURO.0500-19.2020. <https://doi.org/10.1523/ENEURO.0500-19.2020>
- LeBlanc, J. J., DeGregorio, G., Centofante, E., Vogel-Farley, V. K., Barnes, K., Kaufmann, W. E., Fagiolini, M., & Nelson, C. A. (2015). Visual evoked potentials detect cortical processing deficits in Rett syndrome: VEP in Rett syndrome. *Annals of Neurology*, 78(5), 775–786. <https://doi.org/10.1002/ana.24513>
- Lee, L.-J., Tsytsarev, V., & Erzurumlu, R. S. (2017). Structural and functional differences in the barrel cortex of *Mecp2* null mice. *Journal of Comparative Neurology*, 525(18), 3951–3961. <https://doi.org/10.1002/cne.24315>
- Lewis, J. D., Meehan, R. R., Henzel, W. J., Maurer-Fogy, I., Jeppesen, P., Klein, F., & Bird, A. (1992). Purification, sequence, and cellular localization of a novel chromosomal protein that binds to methylated DNA. *Cell*, 69(6), 905–914. [https://doi.org/10.1016/0092-8674\(92\)90610-O](https://doi.org/10.1016/0092-8674(92)90610-O)
- Linhoff, M. W., Garg, S. K., & Mandel, G. (2015). A high-resolution imaging approach to investigate chromatin architecture in complex tissues. *Cell*, 163(1), 246–255. <https://doi.org/10.1016/j.cell.2015.09.002>
- Lo, F.-S., Blue, M. E., & Erzurumlu, R. S. (2016). Enhancement of post-synaptic GABA<sub>A</sub> and extrasynaptic NMDA receptor-mediated responses in the barrel cortex of *Mecp2* -null mice. *Journal of Neurophysiology*, 115(3), 1298–1306. <https://doi.org/10.1152/jn.00944.2015>
- Lonstein, J. S., Lévy, F., & Fleming, A. S. (2015). Common and divergent psychobiological mechanisms underlying maternal behaviors in non-human and human mammals. *Hormones and Behavior*, 73, 156–185. <https://doi.org/10.1016/j.yhbeh.2015.06.011>
- Lonstein, J. S., & Stern, J. M. (1999). Effects of unilateral suckling on nursing behavior and c-fos activity in the caudal periaqueductal gray in rats. *Developmental Psychobiology*, 35(4), 264–275. [https://doi.org/10.1002/\(SICI\)1098-2302\(199912\)35:4<264::AID-DEV2>3.0.CO;2-U](https://doi.org/10.1002/(SICI)1098-2302(199912)35:4<264::AID-DEV2>3.0.CO;2-U)
- Lyst, M. J., & Bird, A. (2015). Rett syndrome: A complex disorder with simple roots. *Nature Reviews Genetics*, 16(5), 261–275. <https://doi.org/10.1038/nrg3897>
- Lyst, M. J., Ekiert, R., Ebert, D. H., Merusi, C., Nowak, J., Selfridge, J., Guy, J., Kastan, N. R., Robinson, N. D., de Lima Alves, F., Rappsilber, J., Greenberg, M. E., & Bird, A. (2013). Rett syndrome mutations abolish the interaction of MeCP2 with the NCoR/SMRT co-repressor. *Nature Neuroscience*, 16(7), 898–902. <https://doi.org/10.1038/nn.3434>
- Maier, R. A. (1963). Maternal behavior in the domestic hen: The role of physical contact. *Journal of Comparative and Physiological Psychology*, 56(2), 357–361. <https://doi.org/10.1037/h0045307>
- Martínez de Paz, A., Khajavi, L., Martin, H., Claveria-Gimeno, R., Tom Dieck, S., Cheema, M. S., Sanchez-Mut, J. V., Moksa, M. M., Carles, A., Brodie, N. I., Sheikh, T. I., Freeman, M. E., Petrotchenko, E. V., Borchers, C. H., Schuman, E. M., Zytynicki, M., Velazquez-Campoy, A., Abian, O., Hirst, M., ... Ausió, J. (2019). MeCP2-E1 isoform is a dynamically expressed, weakly DNA-bound protein with different protein and DNA interactions compared to MeCP2-E2. *Epigenetics & Chromatin*, 12(1), 63. <https://doi.org/10.1186/s13072-019-0298-1>
- Meehan, R., Lewis, J. D., & Bird, A. P. (1992). Characterization of MeCP2, a vertebrate DNA binding protein with affinity for methylated DNA. *Nucleic Acids Research*, 20(19), 5085–5092. <https://doi.org/10.1093/nar/20.19.5085>
- Meng, X., Wang, W., Lu, H., He, L., Chen, W., Chao, E. S., Fiorotto, M. L., Tang, B., Herrera, J. A., Seymour, M. L., Neul, J. L., Pereira, F. A., Tang, J., Xue, M., & Zoghbi, H. Y. (2016). Manipulations of MeCP2 in glutamatergic neurons highlight their contributions to Rett and other neurological disorders. *eLife*, 5, e14199. <https://doi.org/10.7554/eLife.14199>
- Merbler, A. M., Byiers, B. J., Hoch, J., Dimian, A. C., Barney, C. C., Feyma, T. J., Beisang, A. A., Bartolomucci, A., & Symons, F. J. (2020). Preliminary evidence that resting state heart rate variability predicts reactivity to tactile stimuli in Rett syndrome. *Journal of Child Neurology*, 35(1), 42–48. <https://doi.org/10.1177/0883073819875915>
- Mitra, S., & Bult-Ito, A. (2021). Bidirectional behavioral selection in mice: A novel pre-clinical approach to examining compulsivity. *Frontiers in Psychiatry*, 12, 716619. <https://doi.org/10.3389/fpsy.2021.716619>
- Miyata, S., & Kitagawa, H. (2017). Formation and remodeling of the brain extracellular matrix in neural plasticity: Roles of chondroitin sulfate and hyaluronan. *Biochimica et Biophysica Acta*, 1861(10), 2420–2434. <https://doi.org/10.1016/j.bbagen.2017.06.010>
- Morello, N., Schina, R., Pilotto, F., Phillips, M., Melani, R., Plicato, O., Pizzorusso, T., Pozzo-Miller, L., & Giustetto, M. (2018). Loss of *Mecp2* causes atypical synaptic and molecular plasticity of parvalbumin-expressing interneurons reflecting Rett syndrome-like sensorimotor defects. *eNeuro*, 5(5), ENEURO.0086-18.2018. <https://doi.org/10.1523/ENEURO.0086-18.2018>
- Morgan, H. D., Fleming, A. S., & Stern, J. M. (1992). Somatosensory control of the onset and retention of maternal responsiveness in primiparous Sprague-Dawley rats. *Physiology & Behavior*, 51(3), 549–555. [https://doi.org/10.1016/0031-9384\(92\)90178-5](https://doi.org/10.1016/0031-9384(92)90178-5)
- Mossner, J. M., Batista-Brito, R., Pant, R., & Cardin, J. A. (2020). Developmental loss of MeCP2 from VIP interneurons impairs cortical function and behavior. *eLife*, 9, e55639. <https://doi.org/10.7554/eLife.55639>
- Nakagawa, F., Schulte, B. A., Wu, J. Y., & Spicer, S. S. (1986). GABAergic neurons of rodent brain correspond partially with those staining for glycoconjugate with terminal N-acetylgalactosamine. *Journal of Neurocytology*, 15(3), 389–396.
- Neul, J. L. (2019). Can Rett syndrome be diagnosed before regression? *Neuroscience & Biobehavioral Reviews*, 104, 158–159. <https://doi.org/10.1016/j.neubiorev.2019.07.005>

- Nomura, Y. (2005). Early behavior characteristics and sleep disturbance in Rett syndrome. *Brain and Development*, 27, S35–S42. <https://doi.org/10.1016/j.braindev.2005.03.017>
- Noutel, J., Hong, Y. K., Leu, B., Kang, E., & Chen, C. (2011). Experience-dependent retinogeniculate synapse remodeling is abnormal in MeCP2-deficient mice. *Neuron*, 70(1), 35–42. <https://doi.org/10.1016/j.neuron.2011.03.001>
- Orefice, L. L., Mosko, J. R., Morency, D. T., Wells, M. F., Tasnim, A., Mozeika, S. M., Ye, M., Chirila, A. M., Emanuel, A. J., Rankin, G., Fame, R. M., Lehtinen, M. K., Feng, G., & Ginty, D. D. (2019). Targeting peripheral somatosensory neurons to improve tactile-related phenotypes in ASD models. *Cell*, 178(4), 867–886.e24. <https://doi.org/10.1016/j.cell.2019.07.024>
- Orefice, L. L., Zimmerman, A. L., Chirila, A. M., Sleboda, S. J., Head, J. P., & Ginty, D. D. (2016). Peripheral mechanosensory neuron dysfunction underlies tactile and behavioral deficits in mouse models of ASDs. *Cell*, 166(2), 299–313. <https://doi.org/10.1016/j.cell.2016.05.033>
- Orlando, C., Ster, J., Gerber, U., Fawcett, J. W., & Raineteau, O. (2012). Perisynaptic chondroitin sulfate proteoglycans restrict structural plasticity in an integrin-dependent manner. *Journal of Neuroscience*, 32(50), 18009–18017. <https://doi.org/10.1523/JNEUROSCI.2406-12.2012>
- Pacchiarini, N., Fox, K., & Honey, R. C. (2017). Perceptual learning with tactile stimuli in rodents: Shaping the somatosensory system. *Learning & Behavior*, 45(2), 107–114. <https://doi.org/10.3758/s13420-017-0269-y>
- Patrizi, A., Awad, P. N., Chattopadhyaya, B., Li, C., Di Cristo, G., & Fagioli, M. (2020). Accelerated hyper-maturation of parvalbumin circuits in the absence of MeCP2. *Cerebral Cortex*, 30(1), 256–268. <https://doi.org/10.1093/cercor/bhz085>
- Peters, S. U., Gordon, R. L., & Key, A. P. (2015). Induced gamma oscillations differentiate familiar and novel voices in children with MECP2 duplication and Rett syndromes. *Journal of Child Neurology*, 30(2), 145–152. <https://doi.org/10.1177/0883073814530503>
- Picard, N., & Fagioli, M. (2019). MeCP2: An epigenetic regulator of critical periods. *Current Opinion in Neurobiology*, 59, 95–101. <https://doi.org/10.1016/j.conb.2019.04.004>
- Piekarski, D. J., Boivin, J. R., & Wilbrecht, L. (2017). Ovarian hormones organize the maturation of inhibitory neurotransmission in the frontal cortex at puberty onset in female mice. *Current Biology*, 27(12), 1735–1745.e3. <https://doi.org/10.1016/j.cub.2017.05.027>
- Pizzorusso, T., Medini, P., Berardi, N., Chierzi, S., Fawcett, J. W., & Maffei, L. (2002). Reactivation of ocular dominance plasticity in the adult visual cortex. *Science*, 298(5596), 1248–1251. <https://doi.org/10.1126/science.1072699>
- Rais, M., Binder, D. K., Razak, K. A., & Ethell, I. M. (2018). Sensory processing phenotypes in fragile X syndrome. *ASN Neuro*, 10, 175909141880109. <https://doi.org/10.1177/1759091418801092>
- Renthal, W., Boxer, L. D., Hrvatin, S., Li, E., Silberfeld, A., Nagy, M. A., Griffith, E. C., Vierbuchen, T., & Greenberg, M. E. (2018). Characterization of human mosaic Rett syndrome brain tissue by single-nucleus RNA sequencing. *Nature Neuroscience*, 21(12), 1670–1679. <https://doi.org/10.1038/s41593-018-0270-6>
- Rett, A. (1966). ber ein eigenartiges hirnatrophisches syndrom bei hyperammonemia in kindersalter. (On an unusual brain atrophy syndrome with hyperammonemia in childhood.). *Wein Med Wochensh*, 116(723), 26.
- Rodgers, C. C., Nogueira, R., Pil, B. C., Greeman, E. A., Park, J. M., Hong, Y. K., Fusi, S., & Bruno, R. M. (2021). Sensorimotor strategies and neuronal representations for shape discrimination. *Neuron*, 109(14), 2308–2325.e10. <https://doi.org/10.1016/j.neuron.2021.05.019>
- Rosenblatt, J. S. (1967). Nonhormonal basis of maternal behavior in the rat. *Science*, 156(3781), 1512–1513. <https://doi.org/10.1126/science.156.3781.1512>
- Samaco, R. C., McGraw, C. M., Ward, C. S., Sun, Y., Neul, J. L., & Zoghbi, H. Y. (2013). Female Mecp2<sup>+/−</sup> mice display robust behavioral deficits on two different genetic backgrounds providing a framework for pre-clinical studies. *Human Molecular Genetics*, 22(1), 96–109. <https://doi.org/10.1093/hmg/dd5406>
- Santos-Rebouças, C. B., Boy, R., Vianna, E. Q., Gonçalves, A. P., Piergiorgio, R. M., Abdala, B. B., dos Santos, J. M., Calassara, V., Machado, F. B., Medina-Acosta, E., & Pimentel, M. M. G. (2020). Skewed X-chromosome inactivation and compensatory upregulation of escape genes precludes major clinical symptoms in a female with a large Xq deletion. *Frontiers in Genetics*, 11, 101. <https://doi.org/10.3389/fgene.2020.00101>
- Scaramuzza, L., De Rocco, G., Desiato, G., Cobolli Gigli, C., Chiacchiaretta, M., Mirabella, F., Pozzi, D., De Simone, M., Conforti, P., Pagani, M., Benfenati, F., Cesca, F., Bedogni, F., & Landsberger, N. (2021). The enhancement of activity rescues the establishment of Mecp2 null neuronal phenotypes. *EMBO Molecular Medicine*, 13(4), e12433. <https://doi.org/10.15252/emmm.202012433>
- Schaffler, M. D., Elias, L. J., & Abdus-Saboor, I. (2022). Mechanisms of tactile sensory phenotypes in autism: Current understanding and future directions for research. *Current Psychiatry Reports*, 21(12), 134. <https://doi.org/10.1007/s11920-022-01345-0>
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9(7), 671–675. <https://doi.org/10.1038/nmeth.2089>
- Sigal, Y. M., Bae, H., Bogart, L. J., Hensch, T. K., & Zhuang, X. (2019). Structural maturation of cortical perineuronal nets and their perforating synapses revealed by superresolution imaging. *Proceedings of the National Academy of Sciences of the United States of America*, 116(14), 7071–7076. <https://doi.org/10.1073/pnas.1817222116>
- Skene, P. J., Illingworth, R. S., Webb, S., Kerr, A. R. W., James, K. D., Turner, D. J., Andrews, R., & Bird, A. P. (2010). Neuronal MeCP2 is expressed at near histone-octamer levels and globally alters the chromatin state. *Molecular Cell*, 37(4), 457–468. <https://doi.org/10.1016/j.molcel.2010.01.030>
- Stallworth, J. L., Dy, M. E., Buchanan, C. B., Chen, C.-F., Scott, A. E., Glaze, D. G., Lane, J. B., Lieberman, D. N., Oberman, L. M., Skinner, S. A., Tierney, A. E., Cutter, G. R., Percy, A. K., Neul, J. L., & Kaufmann, W. E. (2019). Hand stereotypies: Lessons from the Rett syndrome natural history study. *Neurology*, 92(22), e2594–e2603. <https://doi.org/10.1212/WNL.0000000000007560>
- Stauffer, W., Sheng, H., & Lim, H. N. (2018). EzColocalization: An ImageJ plugin for visualizing and measuring colocalization in cells and organisms. *Scientific Reports*, 8, 15764. <https://doi.org/10.1038/s41598-018-33592-8>
- Stern, J. M. (1996). Somatosensation and maternal care in Norway rats. In *Advances in the study of behavior* (Vol. 25, pp. 243–294). Elsevier. [https://doi.org/10.1016/S0065-3454\(08\)60335-6](https://doi.org/10.1016/S0065-3454(08)60335-6)
- Stern, J. M., & Kolunje, J. M. (1989). Perioral anesthesia disrupts maternal behavior during early lactation in Long-Evans rats. *Behavioral and Neural Biology*, 52(1), 20–38.
- Stern, J. M., & Mackinnon, D. A. (1978). Sensory regulation of maternal behavior in rats: Effects of pup age. *Developmental Psychobiology*, 11(6), 579–586. <https://doi.org/10.1002/dev.420110607>
- Stevenson, P. K., Casenhiser, D. M., Lau, B. Y. B., & Krishnan, K. (2021). Systematic analysis of goal-related movement sequences during maternal behaviour in a female mouse model for Rett syndrome. *European Journal of Neuroscience*, 54(2), 4528–4549. <https://doi.org/10.1111/ejn.15327>
- Su, S.-H., Kao, F.-C., Huang, Y.-B., & Liao, W. (2015). MeCP2 in the rostral striatum maintains local dopamine content critical for psychomotor control. *Journal of Neuroscience*, 35(15), 6209–6220. <https://doi.org/10.1523/JNEUROSCI.4624-14.2015>
- Symons, F. J., Barney, C. C., Byiers, B. J., McAdams, B. D., Foster, S. X. Y. L., Feyma, T. J., Wendelschafer-Crabb, G., & Kennedy, W. R. (2019). A clinical case-control comparison of epidermal innervation density in Rett syndrome. *Brain and Behavior*, 9(5), e01285. <https://doi.org/10.1002/brb3.1285>

- Takei, Y., Zheng, S., Yun, J., Shah, S., Pierson, N., White, J., Schindler, S., Tischbirek, C. H., Yuan, G.-C., & Cai, L. (2021). Single-cell nuclear architecture across cell types in the mouse brain. *Science*, 374(6567), 586–594. <https://doi.org/10.1126/science.abj1966>
- Tao, J., Hu, K., Chang, Q., Wu, H., Sherman, N. E., Martinowich, K., Klose, R. J., Schanen, C., Jaenisch, R., Wang, W., & Sun, Y. E. (2009). Phosphorylation of MeCP2 at Serine 80 regulates its chromatin association and neurological function. *Proceedings of the National Academy of Sciences of the United States of America*, 106(12), 4882–4887. <https://doi.org/10.1073/pnas.0811648106>
- Thompson, E. H., Lensjø, K. K., Wigestrand, M. B., Malthe-Sørensen, A., Hafting, T., & Fyhn, M. (2018). Removal of perineuronal nets disrupts recall of a remote fear memory. *Proceedings of the National Academy of Sciences of the United States of America*, 115, 607–612.
- Ueno, H., Suemitsu, S., Okamoto, M., Matsumoto, Y., & Ishihara, T. (2017). Sensory experience-dependent formation of perineuronal nets and expression of Cat-315 immunoreactive components in the mouse somatosensory cortex. *Neuroscience*, 355, 161–174. <https://doi.org/10.1016/j.neuroscience.2017.04.041>
- Ueno, M., Nakamura, Y., Li, J., Gu, Z., Niehaus, J., Maezawa, M., Crone, S. A., Goulding, M., Baccei, M. L., & Yoshida, Y. (2018). Corticospinal circuits from the sensory and motor cortices differentially regulate skilled movements through distinct spinal interneurons. *Cell Reports*, 23(5), 1286–1300.e7. <https://doi.org/10.1016/j.celrep.2018.03.137>
- Veeraragavan, S., Wan, Y.-W., Connolly, D. R., Hamilton, S. M., Ward, C. S., Soriano, S., Pitcher, M. R., McGraw, C. M., Huang, S. G., Green, J. R., Yuva, L. A., Liang, A. J., Neul, J. L., Yasui, D. H., LaSalle, J. M., Liu, Z., Paylor, R., & Samaco, R. C. (2016). Loss of MeCP2 in the rat models regression, impaired sociability and transcriptional deficits of Rett syndrome. *Human Molecular Genetics*, 25(15), 3284–3302. <https://doi.org/10.1093/hmg/ddw178>
- Wu, H.-P. P., & Dyck, R. H. (2018). Signaling by synaptic zinc is required for whisker-mediated, fine texture discrimination. *Neuroscience*, 369, 242–247. <https://doi.org/10.1016/j.neuroscience.2017.11.020>
- Wu, H.-P. P., Ioffe, J. C., Iverson, M. M., Boon, J. M., & Dyck, R. H. (2013). Novel, whisker-dependent texture discrimination task for mice. *Behavioural Brain Research*, 237, 238–242. <https://doi.org/10.1016/j.bbr.2012.09.044>
- Yagasaki, Y., Miyoshi, G., & Miyata, M. (2018). Experience-dependent MeCP2 expression in the excitatory cells of mouse visual thalamus. *PLoS ONE*, 13(5), e0198268. <https://doi.org/10.1371/journal.pone.0198268>
- Ye, Q., & Miao, Q.-L. (2013). Experience-dependent development of perineuronal nets and chondroitin sulfate proteoglycan receptors in mouse visual cortex. *Matrix Biology*, 32(6), 352–363. <https://doi.org/10.1016/j.matbio.2013.04.001>
- Young, J. I., & Zoghbi, H. Y. (2004). X-chromosome inactivation patterns are unbalanced and affect the phenotypic outcome in a mouse model of Rett syndrome. *American Journal of Human Genetics*, 74(3), 511–520. <https://doi.org/10.1086/382228>
- Yu, Z., Guindani, M., Grieco, S. F., Chen, L., Holmes, T. C., & Xu, X. (2022). Beyond t test and ANOVA: Applications of mixed-effects models for more rigorous statistical analysis in neuroscience research. *Neuron*, 110(1), 21–35. <https://doi.org/10.1016/j.neuron.2021.10.030>
- Zhao, Y.-T., Goffin, D., Johnson, B. S., & Zhou, Z. (2013). Loss of MeCP2 function is associated with distinct gene expression changes in the striatum. *Neurobiology of Disease*, 59, 257–266. <https://doi.org/10.1016/j.nbd.2013.08.001>
- Zhou, Z., Hong, E. J., Cohen, S., Zhao, W., Ho, H. H., Schmidt, L., Chen, W. G., Lin, Y., Savner, E., Griffith, E. C., Hu, L., Steen, J. A. J., Weitz, C. J., & Greenberg, M. E. (2006). Brain-specific phosphorylation of MeCP2 regulates activity-dependent Bdnf transcription, dendritic growth, and spine maturation. *Neuron*, 52(2), 255–269. <https://doi.org/10.1016/j.neuron.2006.09.037>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**FILE S1** Statistical reporting for all generated analyses.

**TABLE S1** Statistical reporting for data presented in Figure 1.

**TABLE S2** Statistical reporting for data presented in Figure 3.

**TABLE S3** Statistical reporting for data presented in Figure 5.

**TABLE S4** Statistical reporting for data presented in Figure 6.

**TABLE S5** Statistical reporting for data presented in Figure 7.

**TABLE S6** Statistical reporting for data presented in Figure 8.

**TABLE S7** Statistical reporting for data presented in Figure 9.

**TABLE S8** Statistical reporting for data presented in Figure 10.

**TABLE A** Mouse lines.

**TABLE B** Antibodies used in the present study.

**TABLE C** Resources used in the present study.

Transparent Science Questionnaire for Authors

**How to cite this article:** Mykins, M., Layo-Carris, D., Dunn, L. R., Skinner, D. W., McBryar, A. H., Perez, S., Shultz, T. R., Willems, A., Lau, B. Y. B., Hong, T., & Krishnan, K. (2023). Wild-type MeCp2 expression coincides with age-dependent sensory phenotypes in a female mouse model for Rett syndrome. *Journal of Neuroscience Research*, 00, 1–23. <https://doi.org/10.1002/jnr.25190>